

Body Weights of Female Mice in the Three-month Feed Study of *Usnea* Lichens

Week ^a	0 ppm			15 ppm			30 ppm			60 ppm			180 ppm			360 ppm		
	N ^b	Mean ± SE ^c	N	Mean ± SE	Pct ^d	N	Mean ± SE	Pct	N	Mean ± SE	Pct	N	Mean ± SE	Pct	N	Mean ± SE	Pct	
0	10	17.4 ± 0.3	10	17.0 ± 0.3		10	16.9 ± 0.3		10	17.7 ± 0.3		10	17.7 ± 0.3		10	17.2 ± 0.2		
1	10	18.4 ± 0.2***	10	18.3 ± 0.2	99.5	10	18.1 ± 0.3	98.4	10	18.6 ± 0.3	101.1	10	18.5 ± 0.2	100.5	10	17.2 ± 0.3***	93.5	
2	10	19.2 ± 0.3***	10	19.9 ± 0.2*	103.6	10	19.2 ± 0.3	100.0	10	19.4 ± 0.3	101.0	10	19.7 ± 0.2	102.6	10	18.1 ± 0.2**	94.3	
3	10	20.0 ± 0.3***	10	20.8 ± 0.3**	104.0	10	19.9 ± 0.3	99.5	10	20.3 ± 0.3	101.5	10	20.5 ± 0.2	102.5	10	18.4 ± 0.2***	92.0	
4	10	20.7 ± 0.3***	10	21.3 ± 0.3	102.9	10	20.5 ± 0.4	99.0	10	21.0 ± 0.4	101.4	10	21.0 ± 0.3	101.4	9	18.9 ± 0.2***	91.3	
5	10	21.6 ± 0.3***	10	22.1 ± 0.3	102.3	10	21.1 ± 0.3	97.7	10	21.4 ± 0.5	99.1	10	21.4 ± 0.3	99.1	9	19.1 ± 0.2***	88.4	
6	10	22.1 ± 0.3***	10	22.6 ± 0.3	102.3	10	21.5 ± 0.3	97.3	10	21.8 ± 0.4	98.6	10	21.7 ± 0.2	98.2	9	19.5 ± 0.3***	88.2	
7	10	22.8 ± 0.5***	10	23.1 ± 0.5	101.3	10	22.3 ± 0.5	97.8	10	22.6 ± 0.4	99.1	10	22.0 ± 0.3	96.5	9	19.6 ± 0.2***	86.0	
8	10	23.3 ± 0.4***	10	23.1 ± 0.5	99.1	10	22.6 ± 0.5	97.0	10	22.9 ± 0.6	98.3	10	22.1 ± 0.2*	94.8	9	20.0 ± 0.3***	85.8	
9	9	23.4 ± 0.4***	10	23.3 ± 0.4	99.6	10	22.6 ± 0.5	96.6	10	22.6 ± 0.4	96.6	10	22.7 ± 0.2	97.0	9	20.3 ± 0.3***	86.8	
10	9	22.8 ± 0.4***	10	23.3 ± 0.4	102.2	10	23.1 ± 0.4	101.3	10	23.1 ± 0.5	101.3	10	23.0 ± 0.2	100.9	9	20.4 ± 0.2***	89.5	
11	9	23.3 ± 0.5***	10	23.9 ± 0.6	102.6	10	23.7 ± 0.6	101.7	10	23.5 ± 0.4	100.9	10	23.5 ± 0.2	100.9	9	20.7 ± 0.2***	88.8	
12	9	24.2 ± 0.6***	10	24.4 ± 0.5	100.8	10	23.8 ± 0.6	98.3	10	24.4 ± 0.6	100.8	10	24.0 ± 0.4	99.2	9	20.8 ± 0.4***	86.0	
13	9	24.9 ± 0.7***	10	24.9 ± 0.5	100.0	10	24.3 ± 0.5	97.6	10	24.8 ± 0.6	99.6	10	24.3 ± 0.4	97.6	9	21.7 ± 0.3***	87.1	
Mean for Weeks																		
1–13		22.0 ± 0.2***		22.6 ± 0.2	102.7		22.0 ± 0.2	100.0		21.8 ± 0.2	99.1		21.6 ± 0.2	98.2		19.6 ± 0.2***	89.1	

^aMeasured after each week of exposure.

^bN = number of animals.

^cBody weight (g) as mean ± standard error. Asterisks denote significant exposure trend (control column) or significant pairwise comparison to control group (Dunnett's test, other columns): p ≤ 0.05 (*); p ≤ 0.01 (**); p ≤ 0.001 (***).

^dMean weight as percentage of control.