

ADME NTP Study K10998 Diisodecyl phthalate

The contractor used the abbreviation DIDP for the test article in the comparison tables.

Sex/Species: male F344 rats.

Vehicle: dermal, absolute ethanol.

CASRN 26761-40-0

Radiolabeled with carbon-14 in the phthalyl moiety; Diisodecyl phthalate, [¹⁴C-U-phthalyl]

Studies Performed:

- Single 30 mg/kg dermal dose to rats with covered dose site and sacrifice 7 days postdose. (n = 3)

Di-isodecyl phthalate is one of nine phthalates that were tested together to determine excretion and tissue distribution after dermal administration. Comparison data for tissue distribution with di(2-ethylhexyl) phthalate (DEHP) is shown in Table 2. The comparison data for all nine phthalates is found in the dimethyl phthalate study S0043.

Note on Accessibility: Persons with disabilities or using assistive technology may find some documents are not fully accessible. For assistance, contact [Central Data Management](#) or use our [contact form](#) and identify the documents/pages for which access is required. We will assist you in accessing the content of the files. NIEHS has helpful information on accessibility.

Table 1. Excretion profile of di-isodecyl phthalate in the urine and the feces after dermal application to the rat*

Time (hr)	% Dose Excreted	
	Urine	Feces
24	--	0.04 ± 0.01
48	--	0.057 ± 0.021
72	--	0.06 ± 0.02
96	--	0.062 ± 0.021
120	--	0.098 ± 0.072
144	--	0.073 ± 0.046
168	--	0.099 ± 0.077
Total	--	0.49 ± 0.246

*Male F-344 rats (200 ± 20 gm) received di-isodecyl phthalate in ethanol dermally (30 mg/kg). The skin was covered with a perforated plastic cap. Data points are the mean ± standard deviation. The percentage of dose excreted represents the fraction of the dose found (as ¹⁴C-equivalent) relative to the total ¹⁴C-equivalent applied.

Table 2. Tissue distribution of di-(2-ethyl hexyl) phthalate and di-isodecyl phthalate after 7 days of dermal exposure

Tissue	% Dose Found ($\bar{X} \pm$ S.D., n = 3)	
	DEHP	DIDP
Brain	0.006 \pm 0.005	0.08 \pm 0.08
Lung	0.005 \pm 0.006	0.015 \pm 0.08
Liver	0.063 \pm 0.017	0.058 \pm 0.032
Spleen	0.001 \pm 0.0003	0.002 \pm 0.002
Small Intestine	0.161 \pm 0.158	0.019 \pm 0.011
Kidney	0.012 \pm 0.001	0.01 \pm 0.01
Testis	0.002 \pm 0.001	0.006 \pm 0.003
Fat	0.066 \pm 0.028	0.149 \pm 0.027
Muscle	1.162 \pm 0.22	0.338 \pm 0.05
Skin	0.298 \pm 0.286	0.095 \pm 0.035
Spinal Cord	0.002 \pm 0.0001	0.034 \pm 0.027
Blood	0.023 \pm 0.001	0.015 \pm 0.003
Skin Area of Application	86.74 \pm 16.98	75.88 \pm 1
Plastic Cap	12.163 \pm 3.48	5.571 \pm 2.77
Total Recovery*	105.8 \pm 16.602	82.73 \pm 12.196

*Total recovery represents the sum of the % dose found in the urine, the feces, the tissues and the plastic cap in 7 days.

*Dose applied was 30 mg/kg.