Study Number: MOG11042 Test Type: MOG - Range Finding Route: Dosing in Feed Species/Strain: Rat/Harlan Sprague Dawley

Study Number:

**Study Gender:** 

**PWG Approval Date:** 

Version:

R02: Reproductive Performance Summary Test Compound: Triphenyl Phosphate CAS Number: 115-86-6 Date Report Requested: 02/19/2021 Time Report Requested: 13:08:58 Lab: Battelle

## MOG11042

Both See web page for date of PWG Approval v1.1.7 Study Number: MOG11042 Test Type: MOG - Range Finding Route: Dosing in Feed Species/Strain: Rat/Harlan Sprague Dawley

F0 Female						
	Treatment Groups (ppm)					
	0	1000	3000	10000	15000	30000
No. Females Mated	22	22	15	22	15	15
No. Females Pregnant	19	19	12	18	11	12
No. Females Littering	15	16	12	15	11	
Percent of Pregnant Females/Mated	86.4	86.4	80.0	81.8	73.3	80.0
Percent of Littered Females/Mated	78.9	84.2	80.0	78.9	73.3	
Percent of Littered Females/Pregnant	93.8	100.0	100.0	100.0	100.0	
Gestational Length	22.2 ± 0.1 (15)	22.3 ± 0.1 (16)	22.3 ± 0.1 (12)	22.1 ± 0.1 (15)	22.4 ± 0.2 (11)	

Date Report Requested: 02/19/2021 Time Report Requested: 13:08:58 Lab: Battelle

## LEGEND

F0 time mated females were vendor provided so endpoints involving number of females paired, and pre-coital interval were not calculated. Pregnancy was defined as evidence of implantation or littering. Vendor's time of confirmation of mating was utilized for calculation of F0 gestational length.

Gestation length in days calculated for sperm positive females that delivered a litter.

Animals that died or were removed from study between mating and littering were excluded from the littered/paired, littered/mated, and littered/pregnant endpoints, resulting in percent values that may not reflect the tallies of females paired, mated, or pregnant.

Statistical analysis performed by Cochran-Armitage (trend) and Fisher Exact (pairwise) 2-sided tests for Percent of Mated Females/Paired; Percent of Littered Females/Paired; Percent of Pregnant Females/Mated; Percent of Littered Females/Mated; Percent of Littered Females/Paired; Percent of Littered; Percent of Litter

Statistical analysis performed by Jonckheere (trend) and Shirley or Dunn (pairwise) tests for the Pre-coital Interval and Gestational Length endpoints if present.

Statistical significance for the control group indicates a significant trend test

Statistical significance for a treatment group indicates a significant pairwise test compared to the vehicle control group

\* Statistically significant at P <= 0.05

\*\* Statistically significant at P <= 0.01

The 30,000 ppm group was terminated due to excessive toxicity on GD12.

Three dams were removed from the control, 1000, and 10000 ppm groups for biological sample collection on GD18, and three dams and their litters were removed from these groups on LD4.

\*\* END OF REPORT \*\*