

Study Number: C11054B-01
Test Type: TOX
Route: Dosing in Water
Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)
Test Compound: Sulfolane
CAS Number: 126-33-0

Date Report Requested: 06/22/2022
Time Report Requested: 08:38:48
Lab: Battelle

Study Number: C11054B-01
Study Sex: Both
PWG Approval Date: See web page for date of PWG Approval
Version: v1.4.1
Stat Version: S

Study Number: C11054B-01

I07: Mean Water Consumption (Interim and Chronic Animals)

Date Report Requested: 06/22/2022

Test Type: TOX

Test Compound: Sulfolane

Time Report Requested: 08:38:48

Route: Dosing in Water

CAS Number: 126-33-0

Lab: Battelle with CRL

Species/Strain: Rat/Harlan Sprague Dawley

F0 Females											
Phase	Days	Treatment Groups (mg/L)									
		0			30			100			
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	
Gestation	6 - 9	27.6 ± 0.6	113.2 ± 2.5	41	28.3 ± 0.6	116.8 ± 2.6	33	27.8 ± 0.5	115.4 ± 2.1	35	
	9 - 12	31.7 ± 1.1	123.4 ± 3.8 *	41	32.2 ± 0.8	125.5 ± 2.7	33	31.9 ± 0.7	125.3 ± 2.8	35	
	12 - 15	32.2 ± 0.6	118.5 ± 2.2	41	33.3 ± 0.7	122.1 ± 2.5	33	32.7 ± 0.8	121.1 ± 3.3	35	
	15 - 18	39.7 ± 0.8	133.1 ± 2.5	41	41.2 ± 1.1	136.8 ± 3.2	33	39.6 ± 0.9	132.7 ± 2.7	34	
	18 - 21	39.1 ± 0.7	116.5 ± 2.4	41	40.5 ± 1.2	118.1 ± 3.1	33	39.6 ± 1.2	118.4 ± 4.0	35	
	6 - 21	34.0 ± 0.6	119.7 ± 2.2	41	35.0 ± 0.8	122.5 ± 2.4	33	34.4 ± 0.8	121.9 ± 2.7	35	
Lactation	1 - 4	45.7 ± 2.7	164.4 ± 8.7 **	38	48.8 ± 3.1	175.6 ± 10.3	31	49.6 ± 4.2	177.1 ± 13.8	32	
	4 - 7	48.5 ± 1.1	168.2 ± 4.1	38	50.6 ± 1.0	176.0 ± 3.2	31	48.0 ± 1.3	167.9 ± 4.3	32	
	7 - 10	57.0 ± 1.3	191.3 ± 4.4	38	63.2 ± 1.1 **	211.8 ± 3.7 *	30	59.3 ± 1.8	201.2 ± 6.2	32	
	10 - 14	63.3 ± 1.7 *	206.5 ± 5.7 **	38	68.7 ± 1.3	224.9 ± 4.3 *	30	70.5 ± 2.0 **	230.5 ± 6.2 **	31	
	1 - 14	54.4 ± 1.3	186.0 ± 4.2 **	38	58.3 ± 1.1 *	200.1 ± 3.8 *	31	58.6 ± 1.7 *	201.0 ± 5.4 **	31	
	14 - 17	71.7 ± 1.9 **	232.4 ± 6.1 **	38	75.4 ± 2.5	247.3 ± 7.9 *	31	75.8 ± 2.2 *	248.2 ± 7.0 *	31	
	17 - 21	90.0 ± 2.3	298.4 ± 8.1	38	94.4 ± 1.8	316.9 ± 6.1	31	90.6 ± 2.5	303.4 ± 8.5	31	
	21 - 24	103.5 ± 3.0 **	354.0 ± 10.7 **	38	105.2 ± 2.1	362.1 ± 6.8	31	106.2 ± 2.7	366.3 ± 10.6	31	
	24 - 28	104.1 ± 3.3	368.4 ± 12.1	38	111.1 ± 3.7	394.2 ± 13.5	31	108.2 ± 3.1	385.9 ± 10.8	31	
	1 - 28	74.4 ± 1.7	254.6 ± 5.9 **	38	78.8 ± 1.2	270.8 ± 4.4	31	77.9 ± 1.9	268.3 ± 6.3	31	

Study Number: C11054B-01**Test Type:** TOX**Route:** Dosing in Water**Species/Strain:** Rat/Harlan Sprague Dawley**I07: Mean Water Consumption (Interim and Chronic Animals)****Test Compound:** Sulfolane**CAS Number:** 126-33-0**Date Report Requested:** 06/22/2022**Time Report Requested:** 08:38:48**Lab:** Battelle with CRL**F0 Females**

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Gestation	6 - 9	27.3 ± 0.5	113.8 ± 2.2	37	27.6 ± 0.7	115.2 ± 2.9	38
	9 - 12	32.0 ± 0.9	126.8 ± 4.0	37	32.5 ± 0.7	129.9 ± 3.0	38
	12 - 15	32.7 ± 0.7	121.5 ± 2.5	37	32.4 ± 0.7	121.8 ± 2.8	38
	15 - 18	40.5 ± 0.8	136.1 ± 2.5	37	41.2 ± 1.1	141.3 ± 3.3	38
	18 - 21	39.5 ± 0.8	117.1 ± 2.2	37	41.2 ± 1.6	126.1 ± 4.4	37
	6 - 21	34.4 ± 0.6	121.8 ± 1.9	37	34.9 ± 0.8	126.0 ± 2.6	38
Lactation	1 - 4	45.5 ± 0.8	166.6 ± 3.3	34	46.0 ± 1.0	172.9 ± 3.5 **	34
	4 - 7	48.5 ± 0.9	170.2 ± 2.9	34	49.4 ± 0.9	178.1 ± 3.0	33
	7 - 10	62.0 ± 1.5	209.5 ± 3.8	34	59.9 ± 1.3	208.0 ± 4.6	34
	10 - 14	69.1 ± 1.1	227.1 ± 3.6 **	34	71.6 ± 2.8	241.8 ± 9.5 **	34
	1 - 14	57.3 ± 0.8	198.1 ± 2.6 **	34	57.9 ± 1.3	205.6 ± 4.4 **	34
	14 - 17	77.8 ± 1.5 **	253.5 ± 4.6 **	34	78.5 ± 2.1 **	261.1 ± 6.3 **	34
	17 - 21	91.1 ± 1.8	303.0 ± 5.6	34	94.7 ± 2.8	321.3 ± 9.1	34
	21 - 24	108.7 ± 2.4	374.2 ± 8.1	34	113.3 ± 2.2 *	396.6 ± 7.1 **	34
	24 - 28	110.3 ± 2.7	392.3 ± 9.7	34	112.5 ± 2.2	404.2 ± 8.4	34
	1 - 28	78.1 ± 1.2	269.3 ± 3.7	34	80.1 ± 1.6	282.1 ± 5.2 **	34

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Males: F1 Immunotox

Phase	Days	Treatment Groups (mg/L)								
		0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	16.0 ± 0.4 **	147.1 ± 2.9	31	16.3 ± 0.4	153.4 ± 2.7	31	16.2 ± 0.3	152.6 ± 2.7	32
	35 - 42	24.4 ± 0.6 **	161.8 ± 3.9	31	24.5 ± 0.5	165.4 ± 2.8	31	24.8 ± 0.4	166.4 ± 2.7	32
	42 - 49	28.6 ± 0.5 *	143.1 ± 2.3	31	27.7 ± 0.6	142.1 ± 2.6	31	28.9 ± 0.7	147.5 ± 3.8	32
	49 - 56	28.8 ± 0.5	115.5 ± 1.7	31	28.2 ± 0.6	116.2 ± 2.0	31	29.9 ± 0.7	122.5 ± 3.5	32
	56 - 63	28.7 ± 0.4	104.7 ± 1.5	30	28.4 ± 0.6	106.9 ± 2.1	30	28.7 ± 0.4	106.7 ± 2.0	30

Study Number: C11054B-01
Test Type: TOX
Route: Dosing in Water
Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)
Test Compound: Sulfolane
CAS Number: 126-33-0

Date Report Requested: 06/22/2022
Time Report Requested: 08:38:48
Lab: Battelle with CRL

F1 Males: F1 Immunotox

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	16.4 ± 0.9	154.9 ± 8.1	31	14.5 ± 0.4 *	145.0 ± 4.0	31
	35 - 42	24.5 ± 0.5	166.2 ± 2.5	31	22.4 ± 0.5 **	159.8 ± 2.3	31
	42 - 49	28.4 ± 0.5	145.6 ± 2.3	31	26.4 ± 0.5 **	142.9 ± 1.9	31
	49 - 56	28.9 ± 0.4	118.7 ± 1.3	31	28.0 ± 0.7	120.8 ± 3.0	31
	56 - 63	29.0 ± 0.4	108.3 ± 1.5	31	27.6 ± 0.4	108.3 ± 1.3	30

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Females: F1 Immunotox

Phase	Days	Treatment Groups (mg/L)								
		0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	13.9 ± 0.4 *	146.2 ± 2.9	16	13.6 ± 0.4	146.7 ± 3.6	18	15.0 ± 1.6	158.8 ± 16.6	17
	35 - 42	20.1 ± 0.6	161.0 ± 4.7	16	19.8 ± 0.4	160.7 ± 2.8	18	20.9 ± 0.5	167.0 ± 3.2	17
	42 - 49	21.8 ± 0.7	143.3 ± 3.9	16	21.0 ± 0.5	138.4 ± 2.8	18	21.7 ± 0.6	142.0 ± 3.7	17
	49 - 56	21.3 ± 0.7	120.2 ± 3.2	16	20.6 ± 0.4	117.3 ± 2.2	18	21.0 ± 0.4	118.4 ± 2.1	17
	56 - 63	21.9 ± 0.7	115.6 ± 3.5	16	20.5 ± 0.5	109.4 ± 2.1	18	20.9 ± 0.4	110.4 ± 2.3	17

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Females: F1 Immunotox

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	13.5 ± 0.5	145.0 ± 4.6	17	12.2 ± 0.4	138.6 ± 4.1	18
	35 - 42	19.6 ± 0.5	160.0 ± 2.7	17	19.2 ± 0.8	166.4 ± 8.8	18
	42 - 49	21.1 ± 0.6	139.7 ± 3.0	17	19.8 ± 0.4	138.5 ± 3.3	18
	49 - 56	21.3 ± 0.6	121.2 ± 3.1	17	20.1 ± 0.4	121.1 ± 2.5	18
	56 - 63	21.8 ± 0.7	116.4 ± 3.6	17	20.6 ± 0.6	115.6 ± 3.1	17

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Males: F1 Biosample Animals

Phase	Days	Treatment Groups (mg/L)								
		0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	17.6 ± 0.5	155.7 ± 8.3	4	17.2 ± 0.7	157.4 ± 12.5	3	18.3 ± 1.4	174.6 ± 14.7	5
	35 - 42	25.2 ± 0.6 *	159.6 ± 7.2	4	24.7 ± 1.5	161.0 ± 16.3	3	22.2 ± 2.1	149.3 ± 14.5	5
	42 - 49	28.8 ± 0.9 *	140.0 ± 6.7	4	27.8 ± 0.8	137.4 ± 9.3	3	28.0 ± 0.9	141.1 ± 4.0	5
	49 - 56	29.2 ± 0.7	114.6 ± 4.6	4	29.2 ± 0.8	116.3 ± 7.3	3	28.8 ± 0.4	116.5 ± 1.2	5
	56 - 63	28.2 ± 0.6	94.5 ± 3.9	4	29.2 ± 1.0	99.2 ± 1.5	3	29.0 ± 0.7	100.3 ± 2.6	5
	63 - 70	28.1 ± 0.6	84.3 ± 2.9	4	29.3 ± 1.3	89.8 ± 5.3	3	28.1 ± 1.0	88.0 ± 3.1	5
	70 - 77	27.2 ± 0.6	75.3 ± 2.9	4	27.7 ± 1.8	78.9 ± 4.7	3	27.3 ± 0.9	79.6 ± 2.5	5
	77 - 84	28.6 ± 0.9	73.8 ± 3.9	4	27.2 ± 1.9	73.7 ± 5.4	3	27.5 ± 0.7	75.1 ± 2.1	5
	84 - 91	27.9 ± 0.9	68.7 ± 3.0	4	27.5 ± 1.8	71.2 ± 5.1	3	26.6 ± 1.2	68.9 ± 2.7	5
	91 - 98	27.4 ± 0.7 *	64.8 ± 2.6	4	27.6 ± 1.8	68.4 ± 2.9	3	25.1 ± 0.6	63.0 ± 1.3	5
	98 - 105	27.9 ± 0.6	63.7 ± 2.9	4	27.6 ± 1.6	66.3 ± 2.0	3	25.8 ± 0.9	63.0 ± 2.5	5
	105 - 112	28.8 ± 2.0	63.6 ± 6.3	4	28.4 ± 2.1	66.1 ± 3.2	3	26.4 ± 1.0	62.5 ± 2.6	5
	112 - 119	26.2 ± 0.6	55.9 ± 2.6	4	27.1 ± 1.6	61.2 ± 2.3	3	25.3 ± 1.1	57.9 ± 2.4	5
	28 - 119	27.0 ± 0.6 *	82.5 ± 3.8	4	27.0 ± 1.0	85.8 ± 3.8	3	26.0 ± 0.7	84.1 ± 2.1	5

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Males: F1 Biosample Animals

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	16.9 ± 0.4	157.4 ± 2.2	4	16.4 ± 0.5	155.8 ± 8.1	4
	35 - 42	23.3 ± 1.0	156.5 ± 5.0	4	19.6 ± 1.4	132.1 ± 6.4	4
	42 - 49	28.6 ± 1.9	149.3 ± 9.7	4	25.4 ± 1.0	133.2 ± 8.8	4
	49 - 56	29.2 ± 1.0	124.2 ± 6.1	4	26.6 ± 1.5	112.6 ± 9.5	4
	56 - 63	27.9 ± 1.3	103.9 ± 6.7	4	26.1 ± 1.0	93.7 ± 6.4	4
	63 - 70	27.4 ± 0.7	91.0 ± 2.8	4	25.6 ± 1.8	81.9 ± 7.1	4
	70 - 77	27.2 ± 0.3	82.0 ± 2.4	4	26.1 ± 1.8	76.8 ± 5.7	4
	77 - 84	26.8 ± 0.7	76.3 ± 3.4	4	26.6 ± 2.1	73.7 ± 5.4	4
	84 - 91	26.8 ± 0.9	72.5 ± 4.2	4	25.6 ± 2.1	67.2 ± 5.0	4
	91 - 98	23.5 ± 1.9	61.9 ± 6.3	4	24.2 ± 1.9	60.5 ± 4.4	4
	98 - 105	25.8 ± 1.0	66.1 ± 4.0	4	25.6 ± 2.4	61.8 ± 5.4	4
	105 - 112	25.4 ± 0.9	62.6 ± 4.0	4	24.9 ± 2.5	58.1 ± 5.4	4
	112 - 119	24.0 ± 1.2	58.1 ± 5.1	4	25.0 ± 2.0	56.7 ± 4.3	4
	28 - 119	25.6 ± 0.9	86.3 ± 4.4	4	24.4 ± 1.3	79.5 ± 4.4	4

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Females: F1 Biosample Animals

Phase	Days	Treatment Groups (mg/L)								
		0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	14.3 ± 1.0	149.0 ± 7.3	3	16.3 ± 1.0	168.9 ± 9.1	4	14.6 ± 3.0	148.9 ± 31.4	3
	35 - 42	19.1 ± 1.1	148.6 ± 6.8	3	20.9 ± 1.3	162.7 ± 10.2	4	22.7 ± 3.4	172.7 ± 24.4	3
	42 - 49	21.1 ± 0.7	136.1 ± 3.8	3	22.7 ± 1.4	144.4 ± 10.3	4	22.8 ± 3.8	142.6 ± 20.9	3
	49 - 56	20.1 ± 1.3	113.7 ± 4.1	3	22.1 ± 1.2	122.1 ± 9.2	4	23.2 ± 2.9	127.2 ± 12.9	3
	56 - 63	21.0 ± 1.4	107.0 ± 5.3	3	23.7 ± 0.8	119.6 ± 6.7	4	22.5 ± 2.3	110.7 ± 8.7	3
	63 - 70	21.1 ± 0.7	99.5 ± 2.2	3	23.5 ± 2.3	110.7 ± 14.1	4	24.7 ± 4.0	111.4 ± 13.4	3
	70 - 77	20.8 ± 1.1	92.5 ± 3.6	3	24.3 ± 1.2	105.7 ± 8.4	4	23.3 ± 3.6	99.0 ± 10.8	3
	77 - 84	21.7 ± 1.7	92.2 ± 6.4	3	24.3 ± 2.3	101.1 ± 14.0	4	23.3 ± 2.5	95.1 ± 7.6	3
	84 - 91	21.9 ± 1.0	89.9 ± 4.4	3	22.3 ± 2.2	90.0 ± 12.4	4	24.0 ± 3.2	95.5 ± 9.5	3
	91 - 98	22.1 ± 1.2	88.6 ± 5.6	3	22.5 ± 1.4	88.4 ± 9.4	4	24.2 ± 3.9	93.3 ± 10.8	3
	98 - 105	21.5 ± 0.6	84.5 ± 2.6	3	24.2 ± 2.2	93.2 ± 13.1	4	24.6 ± 4.5	91.7 ± 13.0	3
	105 - 112	21.6 ± 1.0	83.3 ± 5.2	3	24.8 ± 1.5	92.7 ± 9.5	4	24.5 ± 3.2	89.8 ± 8.6	3
	112 - 119	21.3 ± 1.3	80.5 ± 4.5	3	23.5 ± 1.9	86.4 ± 10.8	4	23.5 ± 2.7	83.8 ± 7.7	3
	28 - 119	20.6 ± 1.0	100.4 ± 4.4	3	22.7 ± 1.5	108.6 ± 10.8	4	23.2 ± 3.1	108.3 ± 11.4	3

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Females: F1 Biosample Animals

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	14.3 ± 1.5	143.8 ± 16.6	2	13.3 ± 0.2	141.7 ± 2.1	3
	35 - 42	17.7 ± 2.5	141.0 ± 14.1	2	17.3 ± 0.3	139.4 ± 4.3	3
	42 - 49	18.7 ± 3.6	125.5 ± 18.0	2	20.7 ± 0.3	137.0 ± 5.7	3
	49 - 56	19.6 ± 1.4	110.9 ± 5.8	2	19.6 ± 0.9	112.2 ± 7.2	3
	56 - 63	20.3 ± 0.6	103.2 ± 0.8	2	19.8 ± 1.0	102.3 ± 6.4	3
	63 - 70	24.0 ± 3.9	115.5 ± 22.5	2	20.0 ± 0.8	95.6 ± 5.2	3
	70 - 77	22.1 ± 0.4	98.6 ± 1.3	2	19.6 ± 1.2	88.7 ± 6.2	3
	77 - 84	22.4 ± 0.1	95.2 ± 2.3	2	19.3 ± 0.4	84.8 ± 2.8	3
	84 - 91	21.7 ± 0.2	89.9 ± 3.6	2	22.0 ± 1.7	93.7 ± 10.1	3
	91 - 98	21.3 ± 0.6	85.9 ± 5.4	2	20.7 ± 0.8	84.2 ± 4.0	3
	98 - 105	23.0 ± 0.5	91.4 ± 5.9	2	20.2 ± 0.9	79.9 ± 4.6	3
	105 - 112	21.7 ± 0.2	82.9 ± 3.2	2	19.6 ± 0.8	76.6 ± 1.9	3
	112 - 119	22.2 ± 1.1	82.0 ± 5.6	2	19.8 ± 0.9	75.8 ± 2.8	3
	28 - 119	20.7 ± 0.3	101.1 ± 1.6	2	19.3 ± 0.3	96.0 ± 3.2	3

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Males: F1 Interim Animals

Phase	Days	Treatment Groups (mg/L)								
		0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	18.2 ± 0.7	160.5 ± 6.2	6	16.7 ± 0.5	157.1 ± 6.1	5	18.0 ± 1.2	167.9 ± 13.1	6
	35 - 42	26.0 ± 0.8 **	163.5 ± 4.4 **	6	24.0 ± 0.9	160.2 ± 7.5	5	22.4 ± 1.7	149.1 ± 12.3	6
	42 - 49	29.5 ± 0.8 *	142.1 ± 3.9	6	27.4 ± 0.5	138.7 ± 3.8	5	27.6 ± 0.8	139.6 ± 5.1	6
	49 - 56	29.8 ± 0.7 *	116.5 ± 2.6	6	28.9 ± 0.6	117.9 ± 3.1	5	28.6 ± 0.4	116.4 ± 2.4	6
	56 - 63	28.9 ± 0.6 *	96.5 ± 2.0	6	28.7 ± 0.7	99.9 ± 1.7	5	28.8 ± 0.6	100.4 ± 2.6	6
	63 - 70	28.1 ± 0.5	84.5 ± 1.3	6	28.7 ± 0.9	89.4 ± 2.7	5	27.8 ± 0.9	87.3 ± 3.0	6
	70 - 77	26.6 ± 1.1	74.1 ± 2.6	6	27.8 ± 1.1	80.3 ± 2.8	5	27.0 ± 0.8	78.7 ± 2.5	6
	77 - 84	28.4 ± 0.9	74.7 ± 2.2	6	26.9 ± 1.1	73.3 ± 3.0	5	26.9 ± 0.8	73.8 ± 2.4	6
	84 - 91	27.4 ± 0.7	69.0 ± 1.5	6	27.2 ± 1.1	70.4 ± 2.9	5	26.4 ± 0.9	68.9 ± 2.4	6
	91 - 98	27.3 ± 0.6 **	65.8 ± 1.2	6	27.5 ± 1.0	68.3 ± 1.6	5	25.3 ± 0.5	63.6 ± 1.0	6
	98 - 105	27.3 ± 0.9	63.2 ± 1.8	6	26.9 ± 1.0	64.6 ± 1.6	5	25.6 ± 0.7	62.6 ± 2.1	6
	105 - 112	27.4 ± 1.7	61.6 ± 4.2	6	27.0 ± 1.5	63.3 ± 2.5	5	25.8 ± 1.0	61.1 ± 2.6	6
	112 - 119	25.6 ± 0.7	55.8 ± 1.5	6	26.4 ± 1.0	59.8 ± 1.6	5	25.3 ± 0.9	58.3 ± 2.1	6
	28 - 119	27.0 ± 0.6 *	83.3 ± 2.0	6	26.5 ± 0.7	85.0 ± 2.0	5	25.8 ± 0.6	83.5 ± 2.2	6

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Males: F1 Interim Animals

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	17.1 ± 0.4	153.9 ± 2.7	5	16.2 ± 0.4	155.7 ± 5.3	5
	35 - 42	23.6 ± 0.8	152.0 ± 5.2	5	19.8 ± 1.1 **	135.4 ± 6.7 **	5
	42 - 49	28.9 ± 1.5	143.7 ± 6.7	5	24.8 ± 1.0 **	131.7 ± 6.4	5
	49 - 56	29.0 ± 0.8	118.1 ± 4.3	5	26.0 ± 1.3	111.8 ± 6.6	5
	56 - 63	28.0 ± 1.0	99.8 ± 4.6	5	25.7 ± 0.9	93.8 ± 4.6	5
	63 - 70	27.9 ± 0.7	89.0 ± 1.3	5	25.1 ± 1.5	82.1 ± 5.3	5
	70 - 77	27.2 ± 0.2	79.3 ± 1.0	5	25.4 ± 1.6	76.6 ± 4.7	5
	77 - 84	27.1 ± 0.6	74.3 ± 1.0	5	25.5 ± 1.9	72.6 ± 4.8	5
	84 - 91	27.1 ± 0.8	70.6 ± 1.5	5	24.8 ± 1.8	67.2 ± 4.2	5
	91 - 98	24.3 ± 1.6	61.0 ± 3.5	5	23.6 ± 1.6 *	61.3 ± 3.4	5
	98 - 105	26.2 ± 0.8	64.1 ± 1.4	5	25.0 ± 2.0	62.5 ± 4.3	5
	105 - 112	25.5 ± 0.7	60.5 ± 1.1	5	24.1 ± 2.1	58.3 ± 4.4	5
	112 - 119	24.8 ± 1.2	57.3 ± 2.4	5	24.1 ± 1.8	56.6 ± 3.4	5
	28 - 119	25.9 ± 0.7	83.8 ± 1.7	5	23.8 ± 1.2	79.9 ± 3.5	5

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Females: F1 Interim Animals

Phase	Days	Treatment Groups (mg/L)								
		0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	14.3 ± 1.0	151.9 ± 8.2	3	16.3 ± 1.0	171.1 ± 8.6	4	14.3 ± 2.1	147.4 ± 21.6	4
	35 - 42	19.1 ± 1.1	151.0 ± 6.6	3	20.9 ± 1.3	167.3 ± 8.9	4	21.7 ± 2.6	168.3 ± 17.8	4
	42 - 49	21.1 ± 0.7	137.5 ± 3.2	3	22.7 ± 1.4	149.6 ± 8.8	4	22.0 ± 2.8	139.9 ± 14.9	4
	49 - 56	20.1 ± 1.3	114.8 ± 4.0	3	22.1 ± 1.2	126.1 ± 7.7	4	22.6 ± 2.1	125.2 ± 9.3	4
	56 - 63	21.0 ± 1.4	107.8 ± 5.0	3	23.7 ± 0.8	122.0 ± 4.7	4	22.0 ± 1.7	109.3 ± 6.2	4
	63 - 70	21.1 ± 0.7	99.8 ± 1.2	3	23.5 ± 2.3	110.7 ± 11.1	4	23.8 ± 3.0	108.1 ± 10.1	4
	70 - 77	20.8 ± 1.1	92.9 ± 2.9	3	24.3 ± 1.2	107.0 ± 5.7	4	22.7 ± 2.6	97.2 ± 7.8	4
	77 - 84	21.7 ± 1.7	93.2 ± 6.6	3	24.3 ± 2.3	103.4 ± 11.6	4	23.0 ± 1.8	94.4 ± 5.6	4
	84 - 91	21.9 ± 1.0	90.8 ± 4.9	3	22.3 ± 2.2	91.9 ± 10.2	4	23.1 ± 2.4	92.0 ± 7.8	4
	91 - 98	22.1 ± 1.2	89.0 ± 5.7	3	22.5 ± 1.4	85.4 ± 7.8	4	23.5 ± 2.9	90.6 ± 8.1	4
	98 - 105	21.5 ± 0.6	85.3 ± 2.3	3	24.2 ± 2.2	90.2 ± 11.5	4	23.6 ± 3.3	88.8 ± 9.7	4
	105 - 112	21.6 ± 1.0	84.0 ± 4.9	3	24.8 ± 1.5	94.5 ± 6.9	4	24.1 ± 2.3	88.8 ± 6.2	4
	112 - 119	21.3 ± 1.3	80.9 ± 4.4	3	23.5 ± 1.9	88.0 ± 8.3	4	23.3 ± 1.9	83.6 ± 5.5	4
	28 - 119	20.6 ± 1.0	101.2 ± 4.3	3	22.7 ± 1.5	109.7 ± 8.5	4	22.5 ± 2.3	106.1 ± 8.5	4

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Females: F1 Interim Animals

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	14.6 ± 0.9	147.2 ± 9.5	3	14.1 ± 0.8	117.7 ± 27.4	4
	35 - 42	19.8 ± 2.5	154.4 ± 18.4	3	20.5 ± 3.2	165.7 ± 23.2	4
	42 - 49	21.4 ± 3.5	138.4 ± 18.9	3	23.8 ± 3.1	157.2 ± 17.2	4
	49 - 56	21.5 ± 2.1	119.8 ± 10.9	3	20.4 ± 1.0	120.8 ± 7.8	4
	56 - 63	22.4 ± 2.1	111.9 ± 9.5	3	21.7 ± 2.1	116.1 ± 12.3	4
	63 - 70	23.5 ± 2.3	110.4 ± 13.4	3	23.7 ± 3.8	114.8 ± 16.5	4
	70 - 77	23.3 ± 1.2	101.9 ± 3.8	3	22.7 ± 3.2	103.6 ± 12.5	4
	77 - 84	23.3 ± 0.8	96.8 ± 2.3	3	22.8 ± 3.4	99.9 ± 12.7	4
	84 - 91	21.6 ± 0.1	87.9 ± 2.0	3	24.5 ± 2.8	105.4 ± 11.7	4
	91 - 98	22.4 ± 1.2	88.5 ± 4.3	3	25.0 ± 4.4	102.6 ± 15.7	4
	98 - 105	22.7 ± 0.4	87.5 ± 3.3	3	22.6 ± 2.5	91.3 ± 8.7	4
	105 - 112	23.5 ± 1.9	87.9 ± 5.9	3	23.4 ± 3.8	92.4 ± 13.0	4
	112 - 119	21.5 ± 0.9	79.0 ± 4.2	3	22.5 ± 2.8	87.6 ± 9.4	4
	28 - 119	21.7 ± 1.0	103.8 ± 3.8	3	22.1 ± 2.8	103.8 ± 5.4	4

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Males: F1 Core Animals

Phase	Days	Treatment Groups (mg/L)								
		0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	17.2 ± 0.5	156.4 ± 4.8	27	15.9 ± 0.4	149.9 ± 3.6	25	15.4 ± 0.4 *	143.3 ± 3.0	26
	35 - 42	25.8 ± 0.4 **	167.7 ± 2.5 *	27	23.6 ± 0.6 *	159.5 ± 2.6	25	23.5 ± 0.6 *	157.9 ± 3.6	26
	42 - 49	29.0 ± 0.5 *	144.2 ± 3.0	27	28.6 ± 1.0	147.8 ± 4.8	25	27.2 ± 0.7	138.8 ± 3.1	25
	49 - 56	29.6 ± 0.5 *	119.1 ± 1.8	27	29.1 ± 0.5	121.0 ± 2.0	25	28.3 ± 0.6	116.6 ± 2.1	26
	56 - 63	28.8 ± 0.4	99.2 ± 1.4	27	28.7 ± 0.4	101.7 ± 1.7	25	28.7 ± 0.8	100.8 ± 2.6	26
	63 - 70	27.9 ± 0.4	86.4 ± 1.2	27	27.6 ± 0.5	87.9 ± 1.5	25	27.4 ± 0.5	86.5 ± 1.3	26
	70 - 77	27.2 ± 0.5	77.7 ± 1.4	27	27.5 ± 0.5	80.9 ± 1.3	25	27.0 ± 0.5	78.7 ± 1.1	26
	77 - 84	27.1 ± 0.5	72.9 ± 1.2 *	27	26.8 ± 0.5	74.2 ± 1.4	25	26.9 ± 0.5	74.0 ± 1.2	26
	84 - 91	26.5 ± 0.4	68.1 ± 1.0	27	25.3 ± 0.7	66.6 ± 1.7	25	26.0 ± 0.6	68.3 ± 1.2	26
	91 - 98	26.4 ± 0.4	65.4 ± 1.0	27	26.1 ± 0.4	66.4 ± 1.2	25	25.8 ± 0.5	65.4 ± 1.0	26
	98 - 105	25.8 ± 0.5	61.9 ± 1.0	27	25.8 ± 0.5	63.3 ± 1.2	25	25.5 ± 0.7	62.4 ± 1.3	26
	105 - 112	25.8 ± 0.5	59.9 ± 0.9	27	26.0 ± 0.4	61.7 ± 0.9	25	26.1 ± 0.6	62.3 ± 1.3	26
	112 - 119	25.8 ± 0.4	58.2 ± 0.8	27	25.7 ± 0.5	59.5 ± 1.0	25	25.6 ± 0.6	59.6 ± 1.3	26
	28 - 119	26.4 ± 0.3	84.0 ± 1.0	27	25.9 ± 0.4	84.7 ± 1.0	25	25.6 ± 0.5	83.5 ± 1.2	26
	28 - 741	27.3 ± 0.4	55.1 ± 1.3	16	26.0 ± 0.7	54.2 ± 1.1	15	26.9 ± 1.0	56.0 ± 2.1	14

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Males: F1 Core Animals

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	16.7 ± 0.6	152.7 ± 4.6	25	15.2 ± 0.4 *	147.4 ± 3.2	26
	35 - 42	24.3 ± 0.6 *	160.1 ± 3.9	25	23.0 ± 0.7 **	159.9 ± 4.7 *	26
	42 - 49	28.7 ± 0.6	143.7 ± 2.8	25	26.8 ± 0.5 *	142.3 ± 2.8	26
	49 - 56	29.0 ± 0.4	117.2 ± 1.6	25	28.1 ± 0.6	120.2 ± 2.2	26
	56 - 63	28.6 ± 0.4	99.2 ± 1.4	25	28.6 ± 0.5	104.0 ± 1.8	26
	63 - 70	27.8 ± 0.4	86.7 ± 1.3	25	27.8 ± 0.5	90.1 ± 1.6	26
	70 - 77	27.8 ± 0.5	80.1 ± 1.2	25	27.6 ± 0.6	82.2 ± 1.7	26
	77 - 84	27.9 ± 0.5	75.5 ± 1.2	25	27.8 ± 0.8	77.4 ± 2.0	26
	84 - 91	26.8 ± 0.5	69.3 ± 1.2	25	26.6 ± 0.6	70.7 ± 1.4	26
	91 - 98	25.7 ± 0.7	64.1 ± 1.7	25	26.4 ± 0.6	67.5 ± 1.5	26
	98 - 105	26.6 ± 0.5	64.1 ± 1.0	25	25.9 ± 0.7	64.2 ± 1.5	26
	105 - 112	26.2 ± 0.5	61.3 ± 1.1	25	26.7 ± 0.8	64.2 ± 1.7	26
	112 - 119	26.0 ± 0.5	59.2 ± 0.9	25	26.5 ± 0.8	62.1 ± 1.7	26
	28 - 119	26.3 ± 0.4	84.3 ± 1.0	25	25.9 ± 0.5	86.0 ± 1.4	26
	28 - 741	26.0 ± 0.7	52.2 ± 1.4	14	26.5 ± 1.0	58.6 ± 3.3	12

Study Number: C11054B-01

I07: Mean Water Consumption (Interim and Chronic Animals)

Date Report Requested: 06/22/2022

Test Type: TOX

Test Compound: Sulfolane

Time Report Requested: 08:38:48

Route: Dosing in Water

CAS Number: 126-33-0

Lab: Battelle with CRL

Species/Strain: Rat/Harlan Sprague Dawley

F1 Females: F1 Core Animals

Phase	Days	Treatment Groups (mg/L)								
		0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	15.2 ± 0.4 **	154.6 ± 3.9 *	14	14.6 ± 0.8	154.9 ± 7.3	14	16.1 ± 1.6	170.0 ± 16.2	13
	35 - 42	20.5 ± 1.1 *	160.8 ± 9.6	14	19.5 ± 0.8	156.3 ± 5.3	14	20.7 ± 0.6	167.9 ± 4.7	13
	42 - 49	21.3 ± 0.7	137.0 ± 4.9	14	22.2 ± 0.7	145.2 ± 3.9	14	21.8 ± 0.5	143.4 ± 4.0	13
	49 - 56	20.8 ± 1.1	119.8 ± 6.9	14	21.4 ± 0.6	120.4 ± 2.7	14	21.2 ± 0.4	119.8 ± 2.6	13
	56 - 63	20.5 ± 0.5	106.3 ± 2.8	14	21.9 ± 0.6	110.4 ± 2.1	14	21.2 ± 0.5	107.6 ± 2.6	13
	63 - 70	20.5 ± 0.6	96.1 ± 2.5	14	21.0 ± 0.4	98.1 ± 1.5	14	20.7 ± 0.4	97.3 ± 1.6	13
	70 - 77	20.8 ± 0.7	91.4 ± 2.6	14	20.8 ± 0.4	92.2 ± 1.6	14	21.3 ± 0.5	94.2 ± 2.1	13
	77 - 84	21.3 ± 0.7	89.5 ± 2.6	14	21.7 ± 0.5	91.5 ± 1.3	14	22.1 ± 0.5	93.1 ± 2.3	13
	84 - 91	21.1 ± 0.3	85.6 ± 1.2	14	21.1 ± 0.4	85.6 ± 1.2	14	21.0 ± 0.6	85.4 ± 2.0	13
	91 - 98	21.2 ± 0.6	83.4 ± 2.0	14	21.5 ± 1.2	85.2 ± 4.2	14	21.6 ± 0.6	85.7 ± 2.1	13
	98 - 105	20.9 ± 0.6	80.7 ± 2.2	14	21.1 ± 0.4	82.3 ± 1.5	14	21.7 ± 0.3	83.9 ± 1.5	13
	105 - 112	21.4 ± 0.9	80.7 ± 3.0	14	21.4 ± 0.5	81.5 ± 1.2	14	21.4 ± 0.6	81.3 ± 2.5	13
	112 - 119	21.4 ± 0.6	79.3 ± 1.8	14	21.1 ± 0.4	78.8 ± 1.5	14	21.5 ± 0.6	80.3 ± 1.9	13
	28 - 119	20.5 ± 0.4	99.1 ± 1.9	14	20.7 ± 0.5	100.5 ± 1.5	14	20.9 ± 0.4	101.7 ± 1.9	13
	28 - 742	23.4 ± 0.3	74.7 ± 1.6	12	24.6 ± 0.5	82.3 ± 1.2	13	24.9 ± 0.6	83.3 ± 3.1	12

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

F1 Females: F1 Core Animals

Phase	Days	Treatment Groups (mg/L)					
		300			1000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	13.8 ± 0.6	148.2 ± 5.6	13	13.1 ± 0.4 **	142.4 ± 3.6	15
	35 - 42	17.8 ± 0.6	146.7 ± 5.2	13	18.0 ± 0.7	150.0 ± 5.0	15
	42 - 49	20.6 ± 0.5	137.9 ± 3.1	13	20.2 ± 0.5	136.9 ± 3.1	15
	49 - 56	20.5 ± 0.5	119.3 ± 2.6	13	19.8 ± 0.5	115.5 ± 2.7	15
	56 - 63	21.0 ± 0.5	109.8 ± 2.1	13	20.9 ± 0.5	109.6 ± 3.0	15
	63 - 70	20.5 ± 0.6	98.6 ± 2.7	13	20.5 ± 0.7	99.5 ± 3.0	15
	70 - 77	20.4 ± 0.5	92.9 ± 2.1	13	20.2 ± 0.5	92.9 ± 2.5	15
	77 - 84	21.3 ± 0.6	92.4 ± 2.8	13	19.8 ± 0.8	87.3 ± 3.2	15
	84 - 91	20.3 ± 0.5	84.4 ± 2.2	13	20.5 ± 0.5	87.9 ± 2.9	15
	91 - 98	20.8 ± 0.7	84.1 ± 2.8	13	21.0 ± 0.5	86.9 ± 2.1	15
	98 - 105	20.9 ± 0.8	82.4 ± 2.7	13	20.4 ± 0.4	82.4 ± 1.2	15
	105 - 112	21.1 ± 0.5	81.6 ± 1.7	13	20.2 ± 0.4	80.5 ± 1.8	15
	112 - 119	22.6 ± 1.0	85.6 ± 4.5	13	20.9 ± 0.4	81.7 ± 1.7	15
	28 - 119	20.1 ± 0.4	99.9 ± 2.1	13	19.6 ± 0.3	99.4 ± 1.6	15
	28 - 742	23.3 ± 0.5	78.3 ± 2.5	11	23.4 ± 0.8	84.7 ± 5.0	12

Study Number: C11054B-01

Test Type: TOX

Route: Dosing in Water

Species/Strain: Rat/Harlan Sprague Dawley

I07: Mean Water Consumption (Interim and Chronic Animals)

Test Compound: Sulfolane

CAS Number: 126-33-0

Date Report Requested: 06/22/2022

Time Report Requested: 08:38:48

Lab: Battelle with CRL

LEGEND

Reported as the mean $\pm$ SEM. N is the number of animals, number of cages for group housed adult animals or number of litters.

F0 females were singly housed. All F1 cohorts were group housed

Water consumption values were excluded when excessive spillage was recorded.

Statistical analysis was performed by Jonckheere (trend) and Shirley or Dunn (pairwise) tests.

Statistical significance for the control group indicates a significant trend test

Statistical significance for a treatment group indicates a significant pairwise test compared to the vehicle control group

* Statistically significant at $P \leq 0.05$

** Statistically significant at $P \leq 0.01$

Consumption is not reported for the non-pregnant animals during gestation and lactation phases

Consumption is not reported for animals during mating

Two water weights were excluded from analysis as outliers: One F1 Immunotox control Male PND 42-49, and one F1 Core Animals 100 mg/L Male PND 56-63, along with the corresponding water weight relative to body weight values.

One female in the 30 mg/L dose group was excluded from the analysis of the PND 28-742 consumption due to a cage change during the interval.

**** END OF REPORT ****