Study Number: R16011
Test Type: Teratology
Route: Oral Gavage

Species/Strain: Rat/Harlan Sprague Dawley

Opecies/Ottaini. Navi lanan Oprague Dawle

Study Gender:

C Number:

**PWG Approval Date** 

**I06: Mean Feed Consumption** 

Test Compound: 2-((1-(4-Phenoxyphenoxy)propan-2-yl)oxy)pyridine

CAS Number: 95737-68-1

R16011

Female

See web page for date of PWG Approval

Date Report Requested: 05/23/2019

Time Report Requested: 12:59:00

Lab: Southern Research

Study Number: R16011

Species/Strain: Rat/Harlan Sprague Dawley

Test Type: Teratology

Route: Oral Gavage

**I06: Mean Feed Consumption** 

Test Compound: 2-((1-(4-Phenoxyphenoxy)propan-2-yl)oxy)pyridine

CAS Number: 95737-68-1

Date Report Requested: 05/23/2019 Time Report Requested: 12:59:00

Lab: Southern Research

## F0 Females

## Treatment Groups (mg/kg/day)

Phase	Litter ID	Days	0				62.5	125			
			Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Gestation	А	3 - 6	18.6 ± 0.3	79.2 ± 1.0	23	19.4 ± 0.3	82.5 ± 0.9	20	18.5 ± 0.3	79.4 ± 1.3	22
		6 - 9	18.4 ± 0.3 **	74.1 ± 1.0 **	23	$18.2 \pm 0.3$	73.5 ± 1.1	20	$17.7 \pm 0.3$	72.2 ± 1.0	22
		9 - 12	$19.3 \pm 0.3$	73.1 ± 0.9	23	$19.5 \pm 0.4$	$74.3 \pm 1.1$	20	$18.9 \pm 0.3$	$72.8 \pm 0.9$	22
		12 - 15	$20.5 \pm 0.4$	$72.7 \pm 0.9$	23	$20.9 \pm 0.4$	$74.6 \pm 1.0$	20	$20.1 \pm 0.5$	72.6 ± 1.4	22
		15 - 18	24.1 ± 0.3 *	77.0 ± 0.8 **	23	$24.5 \pm 0.4$	78.7 ± 1.1	20	$24.1 \pm 0.6$	78.7 ± 1.6	22
		18 - 21	26.1 ± 0.4 **	72.6 ± 0.8 **	23	$27.1 \pm 0.5$	75.0 ± 1.1	20	$26.6 \pm 0.6$	75.3 ± 0.8 *	22
		6 - 21	$21.7 \pm 0.3$	73.6 ± 0.5 *	23	$22.0 \pm 0.3$	$75.0 \pm 0.8$	20	$21.5 \pm 0.3$	$74.3 \pm 0.7$	22

Study Number: R16011

Species/Strain: Rat/Harlan Sprague Dawley

**Test Type:** Teratology **Route:** Oral Gavage

**I06: Mean Feed Consumption** 

Test Compound: 2-((1-(4-Phenoxyphenoxy)propan-2-yl)oxy)pyridine

CAS Number: 95737-68-1

Date Report Requested: 05/23/2019 Time Report Requested: 12:59:00

Lab: Southern Research

## F0 Females

	Litter ID	Days	Treatment Groups (mg/kg/day)							
Phase				250		500				
			Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N		
Gestation	Α	3 - 6	18.6 ± 0.4	78.8 ± 1.6	20	19.0 ± 0.4	80.9 ± 1.4	23		
		6 - 9	$17.7 \pm 0.6$	$71.9 \pm 2.5$	20	16.4 ± 0.4 **	66.8 ± 1.3 **	23		
		9 - 12	$18.7 \pm 0.4$	$72.0 \pm 1.4$	20	$19.0 \pm 0.5$	$73.4 \pm 1.4$	23		
		12 - 15	$20.3 \pm 0.4$	73.6 ± 1.5	20	$21.1 \pm 0.4$	76.4 ± 1.1	23		
		15 - 18	$24.6 \pm 0.6$	80.4 ± 1.5	20	26.2 ± 0.6 *	84.3 ± 1.4 **	23		
		18 - 21	27.6 ± 0.5 *	78.2 ± 1.2 **	20	28.2 ± 0.8 **	78.5 ± 1.7 **	23		
		6 - 21	$21.8 \pm 0.4$	75.2 ± 1.0	20	$22.2 \pm 0.4$	$76.2 \pm 0.9$	23		

Study Number: R16011

**Test Type:** Teratology

**I06: Mean Feed Consumption Test Compound:** 2-((1-(4-Phenoxyphenoxy)propan-2-yl)oxy)pyridine

Route: Oral Gavage CAS Number: 95737-68-1

**Species/Strain:** Rat/Harlan Sprague Dawley

Date Report Requested: 05/23/2019 Time Report Requested: 12:59:00

Lab: Southern Research

## **LEGEND**

Reported as the mean ± SEM. N is the number of animals, number of cages for group housed adult animals or number of litters.

Feed consumpition values were excluded when excessive spillage was recorded.

In multiple breeding/littering studies Litter A is the default designation for the first litter; subsequent litters would be B, C etc.

Statistical analysis performed by Jonckheere (trend) and Shirley or Dunn (pairwise) tests.

Statistical significance for the control group indicates a significant trend test

Statistical significance for a treatment group indicates a significant pairwise test compared to the vehicle control group

- \* Statistically significant at P <= 0.05
- \*\* Statistically significant at P <= 0.01

Consumption is not reported for the non-pregnant animals during gestation and lactation phases

\*\* END OF REPORT \*\*