Test Type: CHRONIC

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 **AVERAGE SEVERITY GRADES[b]** 

**CAS Number:** 15625-89-5

Trimethylolpropane triacrylate

First Dose M/F: 12/13/04 / 12/13/04 Lab: SRI

Time Report Requested: 09:26:54

F1\_M3

C Number: C88133C

**Lock Date:** 10/30/2007

**Cage Range:** ALL

ALL **Date Range:** 

**Reasons For Removal:** 25022 ACCK 25021 TSAC 25020 NATD

25019 MSAC

**Removal Date Range:** ALL

**Treatment Groups:** Include ALL

**Study Gender:** Both

**TDMSE Version:** 2.2.0

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Disposition Summary					
	G.E.	65	65	65	
Animals Initially In Study Early Deaths	65	65	65	00	
Moribund Sacrifice	8	7	10	4	
Natural Death	12	8	11	8	
Survivors	12	· ·	••	J	
Terminal Sacrifice	30	35	29	38	
Animals Examined Microscopically	50	50	50	50	
- Animalo Examinos mioroccopicany					
ALIMENTARY SYSTEM					
Esophagus	(50)	(50)	(50)	(49)	
Inflammation, Chronic Active	, ,	, ,	, ,	1 [2.0]	
Gallbladder	(38)	(41)	(37)	(43)	
Intestine Large, Cecum	(42)	(43)	(40)	(44)	
Edema	, ,	, ,	1 [2.0]	, ,	
Hemorrhage				1 [4.0]	
Necrosis			1 [4.0]		
Thrombosis			1 [3.0]		
Intestine Large, Colon	(44)	(44)	(41)	(46)	
Diverticulum				1 [2.0]	
Intestine Large, Rectum	(45)	(44)	(41)	(47)	
Inflammation, Acute			1 [3.0]		
Intestine Small, Duodenum	(40)	(43)	(40)	(46)	
Edema			1 [2.0]		
Inflammation, Chronic			1 [1.0]		
Intestine Small, Ileum	(42)	(43)	(40)	(45)	
Hemorrhage	1 [3.0]				
Hyperplasia, Lymphoid			1 [2.0]		
Inflammation, Acute		2 [2.0]	2 [2.5]		
Inflammation, Chronic Active		1 [3.0]			
Necrosis	1 [1.0]	1 [1.0]	1 [4.0]		
Thrombosis			1 [2.0]		
Epithelium, Hyperplasia		1 [1.0]			
Muscularis, Hyperplasia		1 [2.0]			
*					

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Intention Coroll Information	(40)	(40)	(40)	(4.4)	
Intestine Small, Jejunum	(40)	(42)	(40)	(44)	
Hemorrhage	4 [0 0]		1 [2.0]		
Hyperplasia, Lymphoid	1 [3.0]		3 [3.0]	4 [0 0]	
Inflammation, Granulomatous		4 [0 0]	4 [4 0]	1 [3.0]	
Inflammation, Acute		1 [2.0]	1 [1.0]		
Inflammation, Chronic Active				1 [2.0]	
Mineralization		1 [1.0]			
Necrosis		2 [1.0]	1 [1.0]		
Muscularis, Hyperplasia		1 [2.0]			
Liver	(50)	(50)	(50)	(50)	
Amyloid Deposition			1 [4.0]		
Angiectasis		1 [4.0]	1 [4.0]	1 [3.0]	
Basophilic Focus	2	2	6	4	
Clear Cell Focus	24	23	18	24 [1.0]	
Congestion	2 [4.0]		1 [4.0]		
Cytoplasmic Alteration			1 [2.0]		
Eosinophilic Focus	18	23	26	19	
Fatty Change	1 [2.0]		2 [2.0]		
Fibrosis		1 [2.0]			
Hematopoietic Cell Proliferation	2 [1.0]	3 [1.0]		3 [1.0]	
Hemorrhage	2 [1.5]		1 [3.0]		
Infarct				1 [4.0]	
Inflammation, Chronic		3 [2.3]		2 [2.0]	
Inflammation, Chronic Active	4 [1.3]	7 [1.3]	6 [1.7]	4 [1.3]	
Mineralization			1 [1.0]	1 [1.0]	
Mixed Cell Focus	2	2	2	1	
Necrosis	1 [4.0]			2 [4.0]	
Necrosis, Focal	11 [2.3]	10 [2.1]	11 [2.3]	6 [3.2]	
Tension Lipidosis	1 -1	2 [2.0]	1	1 [1.0]	
Thrombosis		1 [3.0]	1 [3.0]	2 [1.5]	
Bile Duct, Cyst	1 [1.0]	. []	. []	_[]	
Centrilobular, Necrosis	2 [3.5]		5 [3.4]	1 [3.0]	
Hepatocyte, Vacuolization Cytoplasmic	1 [2.0]	3 [2.3]	2 [2.5]	3 [2.0]	
Kupffer Cell, Hyperplasia	. [=.0]	رد.ن	1 [4.0]	○ [=:0]	
Kupffer Cell, Pigmentation	2 [2.5]	2 [1.5]	4 [2.5]	4 [1.8]	
Oval Cell, Hyperplasia	- [v]	2[0]	1 [2.0]	. []	

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
	(0)	(0)	(4.4)	(0)	
Mesentery	(9)	(8)	(11)	(6)	
Fibrosis			1 [2.0]		
Fat, Necrosis	8 [2.3]	8 [2.1]	8 [2.4]	6 [1.8]	
Oral Mucosa	(0)	(1)	(0)	(0)	
Pancreas	(50)	(49)	(45)	(50)	
Atrophy	2 [1.5]	3 [2.0]	5 [1.8]	3 [2.3]	
Cyst	1 [2.0]		1 [2.0]	1 [2.0]	
Hemorrhage	1 [1.0]				
Hyperplasia, Lymphoid	7 [1.3]	12 [1.6]	19 [1.4]	8 [1.3]	
Infiltration Cellular, Lipocyte				1 [2.0]	
Inflammation, Chronic		1 [2.0]			
Inflammation, Chronic Active				1 [2.0]	
Necrosis				1 [1.0]	
Salivary Glands	(50)	(50)	(50)	(50)	
Atrophy	1 [2.0]		1 [3.0]	3 [2.3]	
Hyperplasia, Lymphoid	36 [1.4]	38 [1.6]	37 [1.7]	40 [1.7]	
Mineralization	2 [1.0]		1 [1.0]		
Pigmentation	1 [1.0]				
Stomach, Forestomach	(50)	(49)	(48)	(50)	
Cyst			1 [2.0]		
Edema	3 [2.0]	2 [1.5]	1 [2.0]		
Erosion	1 [1.0]	1 [1.0]		2 [1.5]	
Hemorrhage				1 [2.0]	
Inflammation		1 [1.0]			
Inflammation, Acute		1 [1.0]			
Inflammation, Chronic	2 [2.0]	1 [2.0]	2 [2.5]	1 [2.0]	
Inflammation, Chronic Active	1 [4.0]	1 [2.0]	1 [3.0]	1 [2.0]	
Necrosis	1 [3.0]	. []	. []	. [=.0]	
Ulcer	9 [2.4]	4 [3.0]	5 [2.2]	5 [3.2]	
Epithelium, Hyperplasia	11 [2.8]	10 [2.2]	8 [2.4]	10 [2.6]	
Stomach, Glandular	(48)	(49)	(44)	(49)	
Congestion	(10)	(10)	( ' ' ')	1 [2.0]	
Cyst	14 [2.0]	19 [2.0]	16 [2.0]	21 [2.0]	
Dysplasia	ا۳ [۲.۷]	10 [2.0]	1 [2.0]	2 : [2.0]	
Erosion	4 [1.0]	3 [1.7]	2 [1.0]	2 [1.5]	
Foreign Body	1 [1.0]	5[1.7]	2 [1.0]	ر۱.۰۱	

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

Route: SKIN APPLICATION
Species/Strain: MICE/B6C3F1

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Hemorrhage	1 [2 0]				
Inflammation, Granulomatous	1 [2.0] 1 [1.0]				
Inflammation, Acute	1 [1.0]	2 [1.5]	1 [2.0]	2 [2.0]	
Inflammation, Acute		2 [1.5] 1 [2.0]	1 [2.0]	2 [2:0]	
Inflammation, Chronic Active		1 [2.0]	2 [2.0]		
Metaplasia		1 [2.0]	2 [2.0]	1 [2 0]	
	2 [4 2]		E [4 O]	1 [3.0] 1 [1.0]	
Metaplasia, Squamous	3 [1.3]	0 [4 7]	5 [1.2]		
Mineralization	1 [1.0]	3 [1.7]	2 [1.5]	8 [1.3]	
Necrosis	4.54.01	2 [1.5]	4 [0.0]	1 [3.0]	
Ulcer	1 [1.0]	0.54.03	4 [2.3]	1 [4.0]	
Glands, Ectasia	5 [1.0]	9 [1.2]	5 [1.2]	6 [1.3]	
Glands, Hyperplasia, Cystic				1 [2.0]	
CARDIOVASCULAR SYSTEM					
Blood Vessel	(1)	(0)	(0)	(1)	
Inflammation, Chronic				1 [2.0]	
Thrombosis				1 [4.0]	
Heart	(50)	(50)	(50)	(50)	
Cardiomyopathy	22 [1.1]	27 [1.4]	28 [1.3]	30 [1.3]	
Inflammation, Acute			3 [2.3]		
Inflammation, Chronic Active			1 [2.0]		
Mineralization		1 [2.0]	3 [1.7]	1 [2.0]	
Thrombosis		2 [3.0]	1 [4.0]	1 [1.0]	
Myocardium, Necrosis	2 [3.5]	2 [0.0]	1 [1.0]	. [0]	
ENDOCRINE SYSTEM					
Adrenal Cortex	(50)	(49)	(47)	(50)	
Accessory Adrenal Cortical Nodule	5 [1.4]	4 [1.8]	1 [2.0]	3 [2.0]	
Degeneration, Hydropic	J [1. <del>4</del> ]	1 [4.0]	ا رکان	ل الح. الح. الح. الح. الح. الح. الح. الح	
Hyperplasia		ı [ <del>4</del> .0]		1 [2.0]	
	14 [4 7]	24 [4 0]	17 [0 4]		
Hypertrophy, Focal	14 [1.7]	24 [1.9]	17 [2.1]	17 [1.8]	
Necrosis	07.10.01	00.10.01	00.14.01	1 [2.0]	
Capsule, Hyperplasia	37 [2.0]	33 [2.2]	36 [1.8]	41 [2.1]	

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Adrenal Medulla	(49)	(49)	(46)	(50)	
Hyperplasia	1 [1.0]	4 [1.3]	3 [1.3]	10 [1.2]	
Islets, Pancreatic	(50)	(49)	(44)	(50)	
Hyperplasia	26 [1.7]	33 [1.9]	21 [1.7]	34 [2.1]	
Parathyroid Gland	(48)	(48)	(48)	(50)	
Amyloid Deposition	,	,	1 [1.0]	,	
Cyst	2 [2.0]	3 [2.0]	3 [2.0]	1 [2.0]	
Pituitary Gland	(50)	(48)	(49)	(50)	
Pars Distalis, Cyst	,	3 [2.0]	4 [2.0]	2 [2.0]	
Pars Distalis, Hyperplasia, Focal		1 [2.0]	1 [2.0]	1 [2.0]	
Thyroid Gland	(50)	(47)	(46)	(48)	
Cyst	()	(,	1 [2.0]	(10)	
Hemorrhage			1 [3.0]		
Inflammation, Chronic		1 [2.0]	. [0.0]		
Follicle, Degeneration, Focal	14 [1.3]	15 [1.8]	20 [1.5]	14 [1.9]	
Follicular Cell, Hyperplasia, Focal	[ • ]	1 [1.0]	1 [2.0]	1 [1.0]	
GENERAL BODY SYSTEM					
Tissue NOS	(2)	(1)	(1)	(0)	
Fibrosis	(2)	(1) 1 [2.0]	(1)	(0)	
Necrosis	1 [2.0] 1 [2.0]	1 [2.0]			
INECIOSIS	1 [2.0]				
GENITAL SYSTEM					
Coagulating Gland	(0)	(0)	(1)	(0)	
Necrosis	. ,		1 [4.0]	• •	
Epididymis	(49)	(50)	(50)	(50)	
Angiectasis	1 [2.0]	, ,	, ,	. ,	
Atrophy			1 [2.0]		
Fibrosis			1 [2.0]		
Granuloma Sperm		2 [2.5]	r -1		
r a a ar ar ar ar	0.54.03		4 4 [4 4]	10 [4 4]	
Inflammation, Chronic	6 [1.3]	1011.31	1411.41	1011. <del>4</del> 1	
Inflammation, Chronic Necrosis	6 [1.3]	16 [1.3] 1 [3.0]	14 [1.4]	10 [1.4]	

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Cyst	2 [2.0]	4 [2.0]	5 [2.0]	6 [2.0]	
Inflammation, Chronic	2 [2.0] 16 [2.1]	4 [2.0] 17 [2.5]	18 [2.1]	18 [2.3]	
Inflammation, Chronic Active	1 [4.0]	17 [2.0]	10 [2.1]	10 [2.5]	
Bilateral, Cyst	1 [2.0]				
Prostate	(50)	(49)	(49)	(50)	
Fibrosis	(30)	1 [3.0]	(43)	(00)	
Infiltration Cellular, Lymphoid	1 [2.0]	1 [0.0]			
Inflammation, Suppurative	1 (2.0 <sub>j</sub>		1 [4.0]	1 [4.0]	
Inflammation, Chronic	21 [1.6]	24 [1.5]	23 [1.4]	26 [1.1]	
Necrosis	21 [1.0]	1 [2.0]	1 [4.0]	2 [3.0]	
Seminal Vesicle	(50)	(50)	(49)	(50)	
Infiltration Cellular, Lymphoid	1 [2.0]	(50)	(43)	(50)	
Inflammation, Suppurative	1 [2.0]		1 [3.0]		
Inflammation, Chronic	8 [1.4]	10 [1.3]	7 [3.6] 7 [1.4]	6 [1.2]	
Inflammation, Chronic Active	0 [1.4]	1 [2.0]	7 [11-7]	0 [1.2]	
Necrosis		1 [1.0]	1 [4.0]		
Epithelium, Hyperplasia		1 [2.0]	1 [4.0]		
Testes	(50)	(50)	(50)	(50)	
Hemorrhage	(00)	1 [2.0]	1 [2.0]	(00)	
Inflammation, Chronic		1 [2.0]	. [2.0]		
Mineralization		. [2.0]	1 [3.0]		
Thrombosis			1 [0.0]	1 [4.0]	
Germinal Epithelium, Atrophy	4 [1.5]	6 [2.0]	5 [1.8]	7 [1.9]	
Interstitial Cell, Hyperplasia	1[1.0]	6 [1.0]	3 [1.0]	, [1.0]	
		0 [1.0]	0 [1.0]		
HEMATOPOIETIC SYSTEM					
Bone Marrow	(50)	(49)	(50)	(50)	
Depletion Cellular			1 [3.0]		
Hemorrhage			1 [3.0]		
Hyperplasia	1 [2.0]	2 [2.0]	1 [2.0]	2 [2.0]	
Infiltration Cellular, Histiocyte				1 [4.0]	
Necrosis	1 [4.0]		1 [4.0]	1 [2.0]	
Lymph Node	(5)	(6)	(2)	(1)	
Hyperplasia, Lymphoid	3 [2.0]		1 [1.0]		

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

Test Type: CHRONIC					
Route: SKIN APPLICATION					
Species/Strain: MICE/B6C3F1					

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Pigmentation	1 [2.0]	4.54.03	1 [2.0]		
Bronchial, Hyperplasia, Lymphoid		1 [4.0]			
Iliac, Hyperplasia, Lymphoid		1 [2.0]			
Mediastinal, Ectasia	1 [2.0]				
Mediastinal, Hyperplasia, Lymphoid	1 [2.0]			1 [2.0]	
Pancreatic, Hyperplasia, Lymphoid		1 [3.0]			
Renal, Hemorrhage		1 [4.0]			
Lymph Node, Mandibular	(48)	(48)	(46)	(46)	
Atrophy		1 [2.0]			
Ectasia	1 [2.0]			1 [2.0]	
Hemorrhage	1 [2.0]				
Hyperplasia, Lymphoid	13 [2.1]	20 [1.7]	14 [1.8]	7 [1.6]	
Inflammation, Granulomatous				1 [1.0]	
Mineralization				1 [1.0]	
Lymph Node, Mesenteric	(48)	(47)	(45)	(50)	
Angiectasis				1 [3.0]	
Atrophy	2 [2.5]	2 [1.5]	1 [1.0]	1 [1.0]	
Congestion	1 [3.0]				
Ectasia		2 [2.0]		2 [2.0]	
Hematopoietic Cell Proliferation		1 [2.0]		1 [2.0]	
Hemorrhage	3 [3.0]	5 [2.2]	7 [3.3]	6 [2.7]	
Hyperplasia, Lymphoid	27 [1.9]	30 [1.9]	28 [2.0]	30 [1.5]	
Infiltration Cellular, Lipocyte			• •	1 [3.0]	
Inflammation, Acute			2 [3.0]	1 [2.0]	
Inflammation, Chronic		1 [2.0]			
Inflammation, Chronic Active		,	1 [3.0]		
Spleen	(49)	(48)	(45)	(49)	
Angiectasis	1 [3.0]	(1-)	(1-)	(1-)	
Fibrosis	1 [2.0]				
Hematopoietic Cell Proliferation	15 [3.4]	16 [2.9]	23 [3.0]	19 [2.8]	
Hyperplasia, Lymphoid	[6]	2 [3.5]	1 [3.0]	2 [3.0]	
Necrosis		1 [2.0]	1 [2.0]	1 [1.0]	
Pigmentation	1 [2.0]	1 [2.0]	2 [1.5]	1 [1.0]	
Lymphoid Follicle, Atrophy	ر اد. ۱	2 [2.5]	1 [2.0]	· [·.•]	
Lymphoid Follicle, Hyperplasia	18 [1.7]	25 [1.7]	17 [1.4]	21 [1.5]	
Thymus	(48)	(48)	(44)	(47)	
inyinuo	(40)	(-10)	(-1-1)	(71)	

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

Route: SKIN APPLICATION
Species/Strain: MICE/B6C3F1

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Cost	0 12 01	14 [2 0]	12 [2 0]	15 [2.0]	
Cyst Hemorrhage	8 [2.0]	14 [2.0]	13 [2.0] 2 [2.0]	15 [2.0]	
Hyperplasia, Lymphoid	3 [1.3]	9 [1.4]	2 [2.0] 6 [1.5]	3 [1.7]	
Epithelial Cell, Hyperplasia	3 [1.3]	9 [1.4] 1 [1.0]	0 [1.5]	3[1.7]	
Epitileliai Geli, Hyperpiasia		1 [1.0]			
INTEGUMENTARY SYSTEM					
Skin	(50)	(50)	(50)	(50)	
Cyst Epithelial Inclusion	1				
Edema				1 [2.0]	
Erosion			1 [2.0]		
Hyperkeratosis	1 [4.0]	1 [1.0]	1 [4.0]	1 [2.0]	
Infiltration Cellular, Melanocyte		1 [1.0]			
Inflammation, Chronic				2 [2.0]	
Inflammation, Chronic Active		1 [4.0]	1 [3.0]		
Mineralization		1 [1.0]			
Necrosis		1 [4.0]			
Ulcer	1 [2.0]				
Control, Fibrosis			1 [1.0]		
Control, Hyperkeratosis	17 [1.3]	24 [1.3]	22 [1.4]	21 [1.3]	
Control, Hyperplasia, Melanocyte			1 [1.0]		
Control, Inflammation, Acute	1 [1.0]				
Control, Inflammation, Chronic	8 [1.0]	10 [1.0]	13 [1.1]	16 [1.0]	
Control, Vacuolization Cytoplasmic			1 [1.0]		
Control, Hair Follicle, Dilatation		1 [1.0]		1 [1.0]	
Control, Subcutaneous Tissue, Hemorrhage		1 [3.0]			
Control Epidermis, Hyperplasia	2 [1.5]		3 [1.0]	3 [1.3]	
Epidermis, Hyperplasia	1 [4.0]				
Epidermis, Site Of Application, Hyperplasia	10 [1.3]	7 [1.9]	15 [1.2]	44 [1.7]	
Hair Follicle, Site Of Application, Dilatation		1 [1.0]	1 [1.0]		
Lip, Inflammation, Acute			1 [2.0]		
Prepuce, Ulcer		1 [4.0]			
Sebaceous Gland, Hyperplasia	1 [4.0]				
Sebaceous Gland, Site Of Application, Hyperplasia		1 [1.0]			

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Cita Of Application Francis				4 [2 0]	
Site Of Application, Erosion Site Of Application, Fibrosis		1 [1.0]		1 [2.0]	
Site Of Application, Fibrosis Site Of Application, Hemorrhage		1 [2.0]			
Site Of Application, Hyperkeratosis	45 [1.8]	50 [2.0]	49 [2.2]	50 [2.2]	
Site Of Application, Hyperplasia, Melanocyte	40 [1.0]	00 [Z.0]	40 [Z.Z]	19 [1.1]	
Site Of Application, Inflammation, Acute	1 [2.0]	2 [2.0]		3 [1.7]	
Site Of Application, Inflammation, Chronic	13 [1.2]	17 [1.1]	26 [1.1]	43 [1.3]	
Site Of Application, Ulcer	10[1.2]	2 [3.0]	20 [1.1]	2 [3.0]	
Site Of Application, Vacuolization Cytoplasmic		2 [0.0]	1 [1.0]	1 [3.0]	
One of Application, Vacabilization bytopiasinic			. [0]	. [6.6]	
MUSCULOSKELETAL SYSTEM					
Bone	(50)	(50)	(50)	(50)	
Osteopetrosis		1 [2.0]			
Cranium, Osteopetrosis	1 [2.0]			1 [2.0]	
Skeletal Muscle	(3)	(2)	(6)	(1)	
Angiectasis			1 [1.0]		
Degeneration	2 [2.5]				
Hemorrhage			1 [3.0]	1 [1.0]	
Inflammation, Chronic		1 [1.0]			
Inflammation, Chronic Active	1 [2.0]				
Necrosis			1 [3.0]		
Capillary, Hyperplasia			1 [2.0]		
NERVOUS SYSTEM					
Brain	(50)	(50)	(50)	(50)	
Congestion	1 [2.0]	1 [2.0]			
Demyelination		1 [2.0]		2 [2.0]	
Gliosis	1 [1.0]			1 [3.0]	
Hemorrhage	4 [1.0]	3 [2.0]	3 [1.0]	5 [1.0]	
Hyperplasia, Lymphoid	1 [1.0]	· -	1 [1.0]	1 [1.0]	
Inflammation, Acute	1 [1.0]				
Inflammation, Chronic	1 [2.0]				
Necrosis	2 [1.5]		3 [1.7]		

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

Lab: SRI

86C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Peripheral Nerve	(2)	(1)	(4)	(1)	
Degeneration	1 [3.0]	1 [1.0]	3 [2.7]	1 [1.0]	
Spinal Cord	(2)	(1)	(4)	(0)	
RESPIRATORY SYSTEM					
Lung	(50)	(50)	(50)	(50)	
Congestion	1 [2.0]		2 [3.5]		
Fibrosis				1 [2.0]	
Hemorrhage	7 [1.9]	5 [1.8]	3 [2.0]	4 [1.3]	
Hyperplasia, Lymphoid	1 [2.0]		1 [2.0]		
Infiltration Cellular, Histiocyte	13 [2.5]	14 [2.4]	9 [1.6]	13 [2.3]	
Inflammation, Suppurative	1 [2.0]	1 [2.0]		1 [1.0]	
Inflammation, Chronic	1 [2.0]			1 [3.0]	
Metaplasia, Osseous	5 [1.0]	1 [1.0]			
Mineralization	1 [1.0]		1 [1.0]	2 [1.0]	
Thrombosis	4 [1.8]	1 [2.0]	3 [2.3]	2 [1.5]	
Alveolar Epithelium, Hyperplasia	6 [2.2]	9 [2.3]	7 [1.6]	5 [2.4]	
Glands, Hyperplasia	1 [2.0]				
Nose	(50)	(50)	(50)	(50)	
Foreign Body		1 [1.0]	2 [2.0]		
Hemorrhage	1 [2.0]				
Inflammation, Chronic	7 [1.6]	11 [1.7]	5 [1.8]	9 [1.6]	
Polyp, Inflammatory	2 [2.0]	1 [3.0]			
Glands, Atrophy				1 [2.0]	
Goblet Cell, Hyperplasia			1 [2.0]		
Respiratory Epithelium, Hyperplasia	1 [1.0]	1 [2.0]		3 [1.3]	
Trachea	(50)	(50)	(50)	(50)	
Hemorrhage				1 [2.0]	
Inflammation, Chronic			1 [2.0]		
Necrosis			1 [2.0]		

#### SPECIAL SENSES SYSTEM

Eye (48)(47)(41) (47)

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Atrophy	1 [4.0]			2 [4.0]	
Cataract	1 [1.0]		1 [1.0]	2 [4.0]	
Inflammation, Chronic	1 [2.0]	1 [1.0]	2 [2.5]	2 [2.0]	
Inflammation, Chronic Active	1 [2.0]	1 [3.0]	2 [2.0]	2 [2.0]	
Cornea, Hyperplasia	1 [2.0]	2 [2.5]	1 [3.0]	2 [2.0]	
Lens, Degeneration	1 [2.0]	1 [4.0]	1 [3.0]	1 [4.0]	
Harderian Gland	(50)	(49)	(49)	(50)	
Hyperplasia, Focal	3 [3.3]	6 [2.7]	3 [2.0]	3 [1.7]	
Inflammation, Chronic	26 [1.3]	23 [1.4]	20 [1.3]	28 [1.4]	
illianimation, Chronic	20 [1.3]	23 [1.4]	20 [1.3]	20 [1.4]	
URINARY SYSTEM					
Kidney	(48)	(46)	(42)	(49)	
Amyloid Deposition			1 [4.0]		
Casts Protein	2 [2.0]	1 [3.0]	1 [2.0]	2 [1.5]	
Cyst	13 [2.0]	13 [2.0]	13 [2.0]	9 [2.0]	
Dilatation				1 [1.0]	
Glomerulosclerosis	1 [2.0]		2 [2.5]		
Hyperplasia, Lymphoid	34 [1.9]	38 [1.7]	31 [1.7]	36 [1.9]	
Infarct	3 [2.7]	3 [1.7]	3 [2.7]	4 [1.8]	
Inflammation, Suppurative	1 [2.0]	1 [3.0]	3 [2.3]	1 [2.0]	
Inflammation, Chronic		• •		1 [1.0]	
Metaplasia, Osseous	9 [1.9]	3 [1.7]	5 [2.0]	3 [2.3]	
Mineralization	17 [1.1]	29 [1.1]	17 [1.1]	31 [1.1]	
Nephropathy	39 [1.7]	41 [2.0]	38 [2.0]	42 [1.9]	
Thrombosis		1 -1	1 [4.0]	1 [1.0]	
Artery, Inflammation, Suppurative	1 [3.0]		[]	F -3	
Papilla, Necrosis	2 [2.0]	2 [2.5]		1 [4.0]	
Pelvis, Dilatation	3 [2.3]	3 [2.0]	1 [2.0]	5 [2.2]	
Renal Tubule, Accumulation, Hyaline Droplet	1 [2.0]	o [=.0]	. [=.0]	÷ []	
Renal Tubule, Dilatation, Focal	16 [1.7]	15 [1.4]	14 [1.6]	22 [1.3]	
Renal Tubule, Dilatation, Diffuse	1 [1.0]	1 [2.0]	1 [2.0]	4 [2.0]	
Renal Tubule, Hyperplasia	. []	. [2.0]	. [2.0]	2 [1.5]	
Renal Tubule, Necrosis	1 [1.0]	1 [2.0]	2 [1.5]	1 [4.0]	
Renal Tubule, Pigmentation	1 [1.0]	ا رک.۷	2 [2.5]	1 [4.0]	

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE MALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Transitional Epithelium, Hyperplasia		1 [2.0]		1 [1.0]	
Urethra	(0)	(1)	(2)	(3)	
Inflammation, Acute	,	. ,	( )	1 [2.0]	
Necrosis				2 [3.0]	
Bulbourethral Gland, Infiltration Cellular			1 [4.0]	2 [4.0]	
Bulbourethral Gland, Necrosis		1 [2.0]	1 [4.0]		
Transitional Epithelium, Hyperplasia				1 [2.0]	
Urinary Bladder	(47)	(48)	(42)	(49)	
Edema	, ,	1 [2.0]	, ,	2 [2.5]	
Hemorrhage	1 [3.0]			2 [2.0]	
Hyperplasia, Lymphoid	18 [1.3]	22 [1.2]	16 [1.3]	19 [1.3]	
Inflammation, Chronic		1 [2.0]		2 [2.5]	
Inflammation, Chronic Active		1 [3.0]		· ·	
Mineralization			1 [1.0]		
Necrosis	1 [2.0]	2 [1.0]	1 [3.0]	1 [2.0]	
Ulcer				1 [4.0]	
Muscularis, Degeneration		1 [2.0]			
Transitional Epithelium, Hyperplasia	1 [1.0]	1 [3.0]		2 [2.5]	

<sup>\*\*\*</sup> END OF MALE \*\*\*

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

Inflammation, Chronic Active

Intestine Small, Duodenum

Ulcer

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

Lab: SRI

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Disposition Summary					
Animals Initially In Study	65	65	65	65	
Early Deaths					
Accidently Killed		1			
Moribund Sacrifice	8	7	8	7	
Natural Death	3	11	12	13	
Survivors					
Natural Death			1		
Terminal Sacrifice	39	31	29	30	
Animals Examined Microscopically	50	50	50	50	
ALIMENTARY SYSTEM					
Esophagus	(50)	(50)	(50)	(50)	
Hemorrhage			1 [1.0]		
Inflammation, Chronic Active				2 [2.5]	
Gallbladder	(46)	(41)	(39)	(40)	
Cyst				1 [2.0]	
Pigmentation				1 [3.0]	
Epithelium, Hyperplasia, Adenomatous, Focal				1 [3.0]	
Intestine Large, Cecum	(47)	(42)	(40)	(42)	
Edema	1 [2.0]			1 [3.0]	
Hemorrhage				1 [1.0]	
Hyperplasia, Lymphoid		1 [3.0]			
Intestine Large, Colon	(47)	(44)	(41)	(44)	
Edema				1 [2.0]	
Fibrosis				1 [1.0]	
Hemorrhage				1 [2.0]	
Intestine Large, Rectum	(48)	(44)	(40)	(43)	
Edema	. ,	1 [2.0]	1 [2.0]	• •	
Hemorrhage				1 [1.0]	
Inflammation, Acute			1 [3.0]		

(47)

(42)

1 [4.0]

(40)

1 [2.0]

(39)

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Fibrosis				1 [1.0]	
Necrosis	3 [1.0]				
Epithelium, Hyperplasia	1 [1.0]	(15)	(42)	(4.5)	
Intestine Small, Ileum	(48)	(42)	(40)	(40)	
Fibrosis				1 [2.0]	
Hyperplasia, Lymphoid			1 [2.0]		
Inflammation, Acute	1 [1.0]	3 [1.7]	1 [1.0]	1 [2.0]	
Necrosis	2 [1.5]				
Epithelium, Hyperplasia			1 [2.0]		
Intestine Small, Jejunum	(47)	(41)	(41)	(40)	
Hemorrhage				1 [1.0]	
Hyperplasia, Lymphoid		1 [1.0]			
Inflammation, Acute				1 [2.0]	
Inflammation, Chronic Active	2 [3.5]				
Necrosis	3 [1.3]	3 [1.3]			
Epithelium, Hyperplasia	1 [2.0]			1 [2.0]	
Liver	(50)	(50)	(50)	(50)	
Angiectasis	2 [3.5]	1 [4.0]	2 [4.0]		
Basophilic Focus	1	3	3		
Clear Cell Focus	3	2	5	6	
Cytoplasmic Alteration, Diffuse	1 [2.0]				
Eosinophilic Focus	15	23	21	21	
Fibrosis				2 [3.0]	
Hematopoietic Cell Proliferation	4 [1.3]	4 [1.5]	10 [1.1]	1 [1.0]	
Hemorrhage	2 [2.0]	,	- 1	2 [2.0]	
Infarct	,	1 [3.0]		,	
Inflammation, Chronic		1 [3.0]		1 [2.0]	
Inflammation, Chronic Active	26 [1.5]	32 [1.4]	24 [1.3]	21 [1.4]	
Mineralization	1 [2.0]	2 [1.0]	[]	[]	
Mixed Cell Focus	. [=.0]	_[]		1	
Necrosis, Focal	10 [2.2]	8 [2.5]	11 [2.4]	7 [2.9]	
Tension Lipidosis	1 [2.0]	0 [2.0]	3 [2.0]	. [=.0]	
Thrombosis	. [=.0]	1 [4.0]	ر د.د.	1 [2.0]	
Centrilobular, Atrophy, Diffuse		ر ۲۰۰۰]	1 [3.0]	1 [2.0]	
Centrilobular, Arrophy, Director Centrilobular, Cytoplasmic Alteration		1 [2.0]	1 [0.0]	ا رح.ن]	
Centrilobular, Oytopiasmic Alteration Centrilobular, Degeneration, Granular, Diffuse		1 [2.0]			

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG
Centrilobular, Necrosis	4 [4 0]	2 (2 0)	4 [4 0]	2 [4 0]
Centrilobular, Necrosis  Centrilobular, Hepatocyte, Vacuolization  Cytoplasmic	1 [4.0]	2 [3.0]	1 [4.0] 1 [1.0]	2 [4.0]
Hepatocyte, Hypertrophy				1 [2.0]
Hepatocyte, Vacuolization Cytoplasmic	1 [2.0]	2 [3.0]	2 [2.0]	1 [2.0]
Hepatocyte, Periportal, Hypertrophy, Diffuse	,	[ <u>-</u> ]	1	1 [2.0]
Kupffer Cell, Hyperplasia	1 [4.0]		1 [2.0]	1 [2.0]
Kupffer Cell, Pigmentation	4 [1.8]	5 [1.2]	10 [1.9]	4 [2.0]
Mesentery	(24)	(25)	(25)	(14)
Fibrosis	, ,	, ,	2 [3.0]	1 [3.0]
Inflammation, Chronic	1 [3.0]			
Inflammation, Chronic Active			1 [3.0]	
Thrombosis	1 [2.0]	1 [3.0]		
Fat, Necrosis	20 [2.0]	23 [2.3]	18 [2.4]	9 [2.2]
Pancreas	(49)	(50)	(46)	(49)
Atrophy	3 [1.7]	3 [2.0]	5 [1.8]	7 [2.1]
Cyst			1 [2.0]	1 [2.0]
Fibrosis				1 [1.0]
Hyperplasia	1 [2.0]			
Hyperplasia, Lymphoid	27 [1.5]	28 [1.4]	26 [1.5]	25 [1.8]
Hyperplasia, Melanocyte				1 [2.0]
Hypertrophy	3 [1.7]	1 [1.0]		2 [1.5]
Inflammation, Acute			1 [2.0]	1 [1.0]
Inflammation, Chronic			1 [1.0]	
Salivary Glands	(50)	(50)	(50)	(49)
Atrophy	1 [2.0]	1 [2.0]	1 [1.0]	
Hyperplasia, Lymphoid	47 [2.0]	38 [1.8]	39 [1.9]	38 [1.9]
Infiltration Cellular, Lymphoid		1 [3.0]		
Mineralization	1 [1.0]	1 [1.0]		
Artery, Hypertrophy			1 [2.0]	
Stomach, Forestomach	(50)	(50)	(47)	(50)
Edema	3 [2.3]		1 [3.0]	1 [2.0]
Erosion	1 [1.0]			
Inflammation, Acute	1 [2.0]			
Inflammation, Chronic		1 [2.0]		1 [3.0]
Mineralization		1 [1.0]		

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
				4.50.03	
Necrosis				1 [2.0]	
Ulcer	4 [2.0]	2 [2.0]	4 [2.5]	1 [2.0]	
Epithelium, Hyperplasia	4 [3.0]	7 [1.7]	9 [2.1]	3 [3.0]	
Stomach, Glandular	(49)	(46)	(43)	(47)	
Cyst	24 [2.0]	24 [2.0]	20 [2.0]	15 [2.0]	
Edema	1 [3.0]				
Erosion	2 [1.5]	1 [2.0]		2 [1.5]	
Fibrosis	1 [2.0]				
Hemorrhage			1 [3.0]		
Infiltration Cellular, Lipocyte			1 [2.0]		
Inflammation, Acute			1 [1.0]		
Inflammation, Chronic Active		2 [1.5]		1 [2.0]	
Metaplasia, Squamous		2 [1.0]	1 [2.0]		
Mineralization	1 [1.0]	2 [1.0]	3 [1.3]	4 [1.0]	
Necrosis	1 [1.0]	1 [1.0]			
Pigmentation			1 [4.0]		
Ulcer	1 [3.0]		1 [3.0]		
Epithelium, Dysplasia	1 [2.0]		. []		
Epithelium, Hyperplasia	1 [2.0]				
Glands, Ectasia	6 [1.3]	7 [1.3]	6 [1.3]	7 [1.3]	
CARDIOVASCULAR SYSTEM					
Heart	(50)	(50)	(50)	(50)	
Cardiomyopathy	31 [1.3]	21 [1.2]	19 [1.4]	19 [1.2]	
Mineralization	0.[0]	1 [1.0]	3 [1.3]	1 [1.0]	
Thrombosis	2 [2.5]	. [5]	2 [1.5]	2 [1.5]	
ENDOCRINE SYSTEM					
Adrenal Cortex	(50)	(50)	(46)	(50)	
Accessory Adrenal Cortical Nodule	4 [2.0]	3 [2.0]	3 [1.0]	5 [1.6]	
Degeneration, Fatty	3 [3.0]	ر د.در	2 [3.5]	0 [0]	
Fibrosis	0 [0.0]		2 [0.0]	1 [2.0]	
Hematopoietic Cell Proliferation	1 [3.0]	2 [2.0]	2 [1.5]	3 [1.7]	
·				3[1.7]	
a - Number of animals examined microscopic	cally at site and number	r of animais with lesio	n		

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Hyperplasia, Focal		2 [1.5]	2 [1.5]	2 [2.0]	
Hypertrophy, Focal	1 [2.0]	1 [2.0]	2 [1.0]	1 [4.0]	
Infiltration Cellular, Lymphoid	1 [2.0]	1 [1.0]		1 [4.0]	
Inflammation, Acute		1 [1.0]	1 [1.0]		
Capsule, Hyperplasia	49 [2.1]	48 [2.1]	45 [2.0]	49 [2.0]	
Adrenal Medulla	(50)	(48)	(45)	(50)	
Hyperplasia	7 [2.0]	2 [2.5]	6 [1.5]	(30)	
Islets, Pancreatic	(49)	(50)	(46)	(49)	
Hyperplasia	6 [1.7]	2 [2.5]	2 [1.0]	6 [1.3]	
Parathyroid Gland	(47)	(47)	(44)	(48)	
Pituitary Gland	(49)	(47)	(49)	(50)	
Thrombosis	(40)	(47)	1 [4.0]	(30)	
Pars Distalis, Angiectasis		3 [1.3]	5 [2.8]		
Pars Distalis, Cyst	1 [2.0]	1 [2.0]	2 [2.0]		
Pars Distalis, Hyperplasia	2 [3.5]	1 [3.0]	1 [3.0]	1 [3.0]	
Pars Distalis, Hyperplasia, Focal	7 [2.0]	5 [2.4]	9 [1.9]	5 [2.2]	
Pars Intermedia, Hyperplasia	7 [2.0]	٥ [٤.٠]	1 [3.0]	J [2.2]	
Pars Intermedia, Hyperplasia, Focal			1 [3.0]	1 [2.0]	
Pars Intermedia, Vacuolization Cytoplasmic,	1 [2.0]			1 [2.0]	
Focal	1 [2.0]				
Rathke's Cleft, Hemorrhage			1 [3.0]		
Thyroid Gland	(49)	(49)	(45)	(48)	
Hyperplasia, Lymphoid	, ,	, ,	, ,	1 [2.0]	
Inflammation, Acute	1 [1.0]				
Inflammation, Chronic Active				1 [1.0]	
Necrosis				1 [1.0]	
Follicle, Degeneration, Focal	29 [1.9]	23 [2.0]	27 [1.7]	20 [2.2]	
Follicular Cell, Hyperplasia		1 [4.0]			
Follicular Cell, Hyperplasia, Focal	4 [1.8]	3 [1.0]	5 [2.0]		
GENERAL BODY SYSTEM		_			
Tissue NOS	(3)	(2)	(4)	(0)	
Fibrosis	(-)	1 [1.0]	1 [3.0]	(-)	
Inflammation, Suppurative	1 [4.0]	F.1.41	[]		
illiamination, Supparative	ı [ <del>-</del> .0]				

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Inflammation, Chronic Active			2 [3.0]		
GENITAL SYSTEM					
Clitoral Gland	(49)	(48)	(49)	(49)	
Inflammation, Chronic	3 [2.0]	2 [2.0]	2 [2.0]	3 [2.3]	
Ovary	(47)	(49)	(45)	(48)	
Angiectasis	4 [2.3]	3 [3.3]	2 [4.0]	,	
Cyst	4 [2.0]	10 [2.0]	4 [2.0]	10 [2.0]	
Hemorrhage	12 [1.8]	15 [1.7]	16 [1.9]	11 [1.9]	
Mineralization	1	1 [4.0]	- 1 - 1	,	
Necrosis		1 [2.0]			
Thrombosis	3 [3.3]	1 [4.0]	1 [4.0]	1 [4.0]	
Bilateral, Cyst	0 [0.0]	1 [2.0]	. [•]	. []	
Uterus	(50)	(50)	(50)	(50)	
Angiectasis	1 [2.0]	3 [4.0]	(00)	(00)	
Cyst	. [2.0]	0[0]	1 [2.0]		
Decidual Reaction		1 [3.0]	. [2.0]		
Edema		2 [3.0]	1 [3.0]		
Hyperplasia, Cystic	48 [3.2]	45 [3.3]	46 [3.5]	46 [3.2]	
Hyperplasia, Lymphoid	10 [0.2]	10 [0.0]	1 [2.0]	10 [0.2]	
Inflammation, Suppurative	1 [4.0]		9 [2.8]	1 [4.0]	
Inflammation, Histiocytic, Focal	1 [ 1.0]		1 [3.0]	. [ 1.0]	
Inflammation, Acute		1 [2.0]	1 [0.0]		
Inflammation, Chronic Active		1 [2.0]	2 [4.0]		
Metaplasia, Osseous	2 [2.0]		2 [4.0]		
Necrosis	2 [3.0]	3 [2.0]	3 [3.0]		
Thrombosis	2 [0.0]	1 [3.0]	3 [3.0]		
Myometrium, Hypertrophy	1 [2.0]	1 [3.0]			
Vagina	(0)	(1)	(1)	(1)	
vagilia	(0)	(1)	(1)	(1)	
HEMATOPOIETIC SYSTEM					
Bone Marrow	(50)	(50)	(49)	(50)	
Hyperplasia	4 [2.3]	1 [4.0]	2 [2.5]	3 [2.3]	

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Pigmentation

Hematopoietic Cell Proliferation

Infiltration Cellular, Lymphoid

Lymphoid Follicle, Atrophy Lymphoid Follicle, Hyperplasia

Hyperplasia, Lymphoid

Fibrosis

Necrosis

Pigmentation

Spleen

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

Lab: SRI

6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Infiltration Cellular, Histiocyte	2 [2.0]		2 [2.0]	1 [1.0]	
Lymph Node	(10)	(7)	(16)	(10)	
Ectasia		2 [3.5]	6 [3.3]	2 [4.0]	
Hematopoietic Cell Proliferation	1 [3.0]				
Hemorrhage		1 [4.0]		2 [4.0]	
Iliac, Ectasia	2 [2.5]			1 [2.0]	
Iliac, Hemorrhage	1 [2.0]				
Iliac, Hyperplasia, Lymphoid	4 [2.5]			1 [2.0]	
Mediastinal, Hemorrhage		1 [4.0]			
Mediastinal, Hyperplasia, Lymphoid		1 [3.0]			
Pancreatic, Hyperplasia	1 [4.0]				
Renal, Hyperplasia, Lymphoid	1 [2.0]				
Lymph Node, Mandibular	(50)	(48)	(48)	(48)	
Atrophy				2 [1.5]	
Hematopoietic Cell Proliferation	1 [3.0]				
Hemorrhage	1 [2.0]	1 [2.0]	1 [1.0]	1 [2.0]	
Hyperplasia, Lymphoid	22 [1.8]	14 [2.0]	9 [1.8]	23 [1.8]	
Lymph Node, Mesenteric	(45)	(46)	(42)	(49)	
Atrophy	2 [2.0]	1 [2.0]	3 [2.0]	1 [1.0]	
Ectasia	1 [2.0]	2 [3.5]	3 [2.0]	1 [4.0]	
Fibrosis		1 [2.0]			
Hematopoietic Cell Proliferation	3 [1.7]			1 [2.0]	
Hemorrhage	4 [2.5]	1 [4.0]	2 [3.0]	1 [2.0]	
Hyperplasia, Histiocytic		1 [3.0]			
Hyperplasia, Lymphoid	29 [2.2]	20 [1.9]	21 [2.2]	32 [1.7]	
Necrosis					
Hyperplasia, Lymphoid	29 [2.2]		21 [2.2]	32 [1.7]	

(49)

1 [2.0]

27 [2.9]

17 [1.9]

24 [1.7]

(44)

27 [2.9]

2 [3.0]

1 [4.0]

19 [1.8]

18 [1.4]

(48)

1 [2.0]

29 [2.7]

4 [3.0]

19 [1.9]

19 [1.6]

1 [2.0]

(48)

31 [2.9]

3 [2.7]

1 [2.0]

22 [1.6]

1 [4.0]

22 [1.6]

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

Route: SKIN APPLICATION
Species/Strain: MICE/B6C3F1

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Thymus	(47)	(49)	(44)	(44)	
Cyst	10 [2.0]	11 [2.0]	12 [2.0]	7 [2.0]	
Hemorrhage	9 [2.0]	15 [2.1]	13 [2.1]	10 [1.7]	
Hyperplasia, Lymphoid	15 [2.2]	19 [2.4]	13 [1.7]	22 [1.9]	
Hyperplasia, Reticulum Cell			1 [2.0]		
INTEGUMENTARY SYSTEM					
Mammary Gland	(50)	(50)	(50)	(50)	
Hyperplasia	11 [1.6]	8 [1.9]	8 [1.8]	11 [1.1]	
Inflammation, Chronic			1 [2.0]		
Skin	(50)	(50)	(50)	(50)	
Cyst Epithelial Inclusion				1 [2.0]	
Edema			1 [4.0]	1 [3.0]	
Hyperkeratosis		1 [2.0]		3 [2.3]	
Hyperplasia, Melanocyte			2 [2.0]	2 [2.0]	
Inflammation, Acute		1 [2.0]			
Inflammation, Chronic			1 [2.0]	3 [2.7]	
Inflammation, Chronic Active			1 [3.0]	1 [4.0]	
Ulcer				4 [4.0]	
Control, Edema	1 [3.0]		3 [2.0]		
Control, Hyperkeratosis	40 [1.5]	40 [1.3]	39 [1.3]	35 [1.4]	
Control, Hyperplasia, Melanocyte			1 [1.0]		
Control, Inflammation, Acute	1 [2.0]		1 [2.0]		
Control, Inflammation, Chronic	28 [1.0]	19 [1.1]	26 [1.1]	27 [1.1]	
Control, Necrosis			1 [1.0]		
Control, Epidermis, Hyperplasia	1 [1.0]		1 [1.0]		
Control, Epidermis, Subcutaneous Tissue, Hyperplasia	1 [1.0]				
Epidermis, Hyperplasia		1 [1.0]	3 [2.0]	4 [3.3]	
Epidermis, Site Of Application, Hyperplasia	7 [1.9]	7 [1.6]	15 [1.5]	34 [1.7]	
Hair Follicle, Dilatation				1 [2.0]	
Hair Follicle, Site Of Application, Dilatation			1 [2.0]	1 [2.0]	
Lip, Inflammation, Chronic				1 [2.0]	
Site Of Application, Erosion	1 [1.0]				

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Site Of Application, Fibrosis		1 [1.0]		1 [2.0]	
Site Of Application, Hyperkeratosis	50 [2.1]	49 [2.1]	50 [2.2]	50 [2.6]	
Site Of Application, Hyperplasia, Melanocyte	1 [1.0]	1 [4.0]	3 [1.7]	33 [1.3]	
Site Of Application, Inflammation, Acute	1 [2.0]	1 [3.0]	2 [2.5]	4 [1.5]	
Site Of Application, Inflammation, Chronic	37 [1.1]	36 [1.2]	43 [1.2]	48 [1.5]	
Site Of Application, Necrosis	[]	22 [=]		1 [1.0]	
Site Of Application, Ulcer			3 [3.3]	3 [3.3]	
MUSCULOSKELETAL SYSTEM					
Bone	(50)	(50)	(50)	(50)	
Fibrosis	8 [2.1]	9 [2.1]	13 [1.8]	11 [1.9]	
Fibrous Osteodystrophy	1 [2.0]				
Fracture				1 [3.0]	
Cranium, Fibrosis	1 [2.0]	2 [2.0]			
Femur, Fibrosis		3 [2.3]	1 [2.0]		
Vertebra, Arthrosis			1 [2.0]		
Skeletal Muscle	(8)	(3)	(4)	(3)	
Degeneration	4 [2.0]			1 [4.0]	
Hemorrhage		1 [3.0]			
NERVOUS SYSTEM					
Brain	(50)	(50)	(50)	(50)	
Compression	4 [3.0]	2 [4.0]	3 [2.3]		
Cyst Epithelial Inclusion				1 [2.0]	
Degeneration	1 [2.0]				
Developmental Malformation			1 [2.0]		
Gliosis	4 [1.5]			1 [1.0]	
Hemorrhage	2 [2.0]	1 [1.0]	2 [1.0]	1 [2.0]	
Hyperplasia, Lymphoid	1 [1.0]	-	3 [1.0]	1 [1.0]	
Inflammation, Chronic	-		-	1 [1.0]	
Necrosis	6 [2.8]	2 [2.0]	4 [2.3]	2 [1.5]	
Thrombosis	1 [2.0]				
Meninges, Fibrosis			1 [1.0]		

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate **CAS Number:** 15625-89-5

Time Report Requested: 09:26:54 First Dose M/F: 12/13/04 / 12/13/04

0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
	1 [2 0]			
(2)		(0)	(1)	
		(0)		
		(0)		
(3)	(2)	(0)		
	1 [2 0]		1 [3.0]	
1 [1 0]	1 [3.0]			
	1 [// 0]			
1 [1.0]	1 [4.0]			
(50)	(50)	(50)	(50)	
	,	,		
	4 [1.8]	4 [1.8]		
	• •			
	9 [2.3]	6 [2.3]	2 [2.5]	
	• •			
1 [2.0]				
	1 [2.0]	2 [1.5]		
1 [1.0]			1 [1.0]	
1 [3.0]				
		2 [2.0]	1 [2.0]	
7 [1.9]	2 [2.5]			
,	,		,	
1 [2.0]	1 [2.0]		1 [2.0]	
6 [1.7]	5 [1.0]	6 [1.5]	5 [1.6]	
(48)	(45)	(43)	(45)	
	( )	( /	()	
			1 [3.0]	
	1 [1.0]		. []	
	(3) 3 [3.3] (3)  1 [1.0] 1 [1.0]  (50) 1 [3.0] 4 [2.3] 1 [2.0] 6 [2.2]  1 [2.0] 1 [1.0]  1 [3.0] 7 [1.9] (50) 1 [2.0]	(3) (2) (2) (3) (2) (3) (2) (3) (2) (2) (3) (2) (2) (2) (2) (2) (2) (2) (2) (2) (2	1 [2.0] (3) (2) (0) 3 [3.3] 1 [2.0] (3) (2) (0)  1 [3.0] 1 [1.0] 1 [1.0] 1 [4.0]  (50) (50) (50) (50) 1 [3.0] 4 [2.3] 4 [1.8] 4 [1.8] 1 [2.0] 6 [2.2] 9 [2.3] 6 [2.3]  1 [2.0] 1 [2.0] 2 [1.5] 1 [1.0] 2 [1.0] 1 [3.0] 2 [1.0] 1 [3.0] 1 [3.0] 2 [2.0] 7 [1.9] 2 [2.5] 1 [2.0] (50) (50) (50) 1 [2.0] 1 [2.0] 6 [1.7] 5 [1.0] 6 [1.5]	(3) (2) (0) (1) (1) (3) (3.3] (2) (0) (1) (1) (1) (3) (2) (0) (1) (1) (1) (1) (1) (1) (1) (1) (1) (1

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

### P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

Lab: SRI

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Inflammation, Chronic Active			1 [2.0]		
Necrosis	1 [4.0]				
Capillary, Hyperplasia	1 [2.0]				
Capillary, Cornea, Hyperplasia			1 [2.0]		
Cornea, Hyperplasia	1 [3.0]	2 [1.0]	1 [1.0]		
Retina, Degeneration	1 [1.0]				
Harderian Gland	(50)	(50)	(48)	(49)	
Hyperplasia	,	,	,	1 [3.0]	
Hyperplasia, Focal	3 [2.3]	4 [1.8]	3 [2.3]		
Inflammation, Acute			1 [3.0]		
Inflammation, Chronic	36 [1.4]	39 [1.4]	34 [1.4]	32 [1.5]	
Necrosis				1 [1.0]	
URINARY SYSTEM					
Kidney	(49)	(46)	(43)	(47)	
Casts Protein	4 [1.3]		2 [1.0]	1 [1.0]	
Cyst	2 [2.0]	1 [2.0]	1 [2.0]		
Glomerulosclerosis				1 [3.0]	
Hyperplasia, Lymphoid	43 [1.8]	38 [1.7]	34 [1.8]	38 [1.8]	
Infarct	5 [2.2]	4 [1.8]	6 [2.0]	3 [1.7]	
Infiltration Cellular, Lymphoid		2 [2.5]			
Metaplasia, Osseous	4 [1.5]		2 [1.5]	3 [1.7]	
Mineralization	7 [1.0]	6 [1.3]	6 [1.0]	5 [1.0]	
Nephropathy	34 [1.4]	29 [1.5]	28 [1.4]	36 [1.3]	
Papilla, Necrosis				1 [2.0]	
Pelvis, Degeneration	1 [2.0]				
Pelvis, Dilatation	1 [3.0]		1 [3.0]	1 [3.0]	
Renal Tubule, Accumulation, Hyaline Droplet	4 [3.5]		1 [2.0]	2 [3.0]	
Renal Tubule, Dilatation, Focal	28 [1.4]	20 [1.3]	19 [1.5]	15 [1.3]	
Renal Tubule, Dilatation, Diffuse		1 [2.0]		2 [2.0]	
Renal Tubule, Hyperplasia	1 [2.0]				
Renal Tubule, Necrosis	1 [2.0]		1 [1.0]		
Renal Tubule, Pigmentation	1 [1.0]	2 [4.0]			

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

1 [2.0]

Transitional Epithelium, Hyperplasia

Test Type: CHRONIC

**Route: SKIN APPLICATION** 

Species/Strain: MICE/B6C3F1

# P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 05/17/2010 AVERAGE SEVERITY GRADES[b]

Trimethylolpropane triacrylate CAS Number: 15625-89-5

Time Report Requested: 09:26:54

First Dose M/F: 12/13/04 / 12/13/04

Lab: SRI

B6C3F1 MICE FEMALE	0 MG/KG	0.3 MG/KG	1.0 MG/KG	3.0 MG/KG	
Urinary Bladder	(48)	(48)	(46)	(48)	
Hyperplasia, Lymphoid	40 [1.6]	29 [1.6]	24 [1.6]	31 [1.5]	
Inflammation, Chronic Active Transitional Epithelium, Hyperplasia		1 [4.0]	1 [1.0]		

\*\*\* END OF REPORT \*\*\*

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)