Test Type: CHRONIC

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Species/Strain: RATS/Wistar Han

Lab: BNW

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01

First Dose M/F: 04/14/08 / 04/14/08

F1_Rev.1_RE

NTP Study Number: C20515

Lock Date: 01/12/2011

Cage Range: ALL

Route: RESPIRATORY EXPOSURE WHOLE BODY

Date Range: ALL

Reasons For Removal: ALL

Removal Date Range: ALL

Include ALL **Treatment Groups:**

Study Gender: Both

TDMSE Version: 3.0.1.1_001

PWG Approval Date: 02/21/2013

Species/Strain: RATS/Wistar Han

Route: RESPIRATORY EXPOSURE WHOLE BODY

Test Type: CHRONIC

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3
Disposition Summary				
Animals Initially In Study	50	50	50	50
Early Deaths				
Moribund Sacrifice	17	12	13	15
Natural Death		4	3	2
Survivors				
Terminal Sacrifice	33	34	34	33
Animals Examined Microscopically	50	50	50	50
ALIMENTARY SYSTEM				
Esophagus	(50)	(50)	(50)	(50)
Intestine Large, Cecum	(50)	(48)	(47)	(49)
Inflammation, Acute		1 (2%)		
Necrosis		1 (2%)		
Intestine Large, Colon	(50)	(49)	(47)	(49)
Inflammation, Acute		1 (2%)		
Necrosis		1 (2%)		
Intestine Large, Rectum	(48)	(47)	(47)	(48)
Intestine Small, Duodenum	(50)	(48)	(47)	(49)
Intestine Small, Ileum	(50)	(48)	(47)	(49)
Intestine Small, Jejunum	(50)	(48)	(47)	(49)
Diverticulum		1 (2%)		
Inflammation, Chronic				1 (2%)
Lymphatic, Angiectasis				1 (2%)
Liver	(50)	(50)	(50)	(50)
Angiectasis	3 (6%)	7 (14%)	8 (16%)	7 (14%)
Basophilic Focus	11 (22%)	20 (40%)	18 (36%)	18 (36%)
Clear Cell Focus	29 (58%)	26 (52%)	17 (34%)	26 (52%)
Congestion		1 (2%)		
Eosinophilic Focus			1 (2%)	
Fatty Change				1 (2%)
Inflammation, Granulomatous		1 (2%)		
Mixed Cell Focus	3 (6%)			1 (2%)
Necrosis	2 (4%)	3 (6%)	1 (2%)	1 (2%)

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Bile Duct, Cyst	1 (2%)	1 (2%)			
Bile Duct, Dilatation	3 (6%)	2 (4%)	2 (4%)	4 (8%)	
Bile Duct, Hyperplasia	1 (2%)	2 (4%)			
Bile Duct, Inflammation, Chronic Active	1 (2%)				
Mesentery	(7)	(7)	(1)	(2)	
Fat, Necrosis	6 (86%)	7 (100%)	1 (100%)		
Oral Mucosa	(1)	(0)	(1)	(0)	
Developmental Malformation	1 (100%)		1 (100%)		
Pancreas	(50)	(50)	(49)	(50)	
Atrophy	1 (2%)	4 (8%)	4 (8%)	5 (10%)	
Basophilic Focus	2 (4%)	4 (8%)	2 (4%)	1 (2%)	
Inflammation, Acute		1 (2%)			
Salivary Glands	(50)	(50)	(50)	(50)	
Atrophy			1 (2%)		
Stomach, Forestomach	(50)	(50)	(50)	(50)	
Hyperplasia, Squamous	1 (2%)			1 (2%)	
Inflammation, Chronic Active	1 (2%)				
Ulcer		1 (2%)		2 (4%)	
Stomach, Glandular	(50)	(50)	(49)	(50)	
Necrosis			1 (2%)		
Tooth	(0)	(1)	(0)	(1)	
Dysplasia				1 (100%)	
Inflammation, Chronic Active		1 (100%)			
CARDIOVASCULAR SYSTEM					
Blood Vessel	(50)	(50)	(50)	(50)	
Heart	(50)	(50)	(50)	(50)	
Cardiomyopathy	28 (56%)	40 (80%)	32 (64%)	33 (66%)	
Necrosis	,	,	1 (2%)	,	
ENDOCRINE SYSTEM					
Adrenal Cortex	(50)	(50)	(49)	(50)	
Accessory Adrenal Cortical Nodule	1 (2%)	` '	` '	,	

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Degeneration, Cystic	2 (4%)	1 (2%)	3 (6%)	2 (4%)	
Hyperplasia	26 (52%)	25 (50%)	28 (57%)	26 (52%)	
Hypertrophy	20 (40%)	21 (42%)	14 (29%)	16 (32%)	
Adrenal Medulla	(50)	(48)	(49)	(50)	
Angiectasis				1 (2%)	
Hyperplasia	2 (4%)	1 (2%)	4 (8%)	1 (2%)	
Islets, Pancreatic	(50)	(50)	(49)	(50)	
Parathyroid Gland	(43)	(43)	(44)	(47)	
Hyperplasia	1 (2%)	,	1 (2%)	1 (2%)	
Pituitary Gland	(49)	(50)	(50)	(50)	
Cyst	1 (2%)	,	,	,	
Pars Distalis, Hyperplasia	14 (29%)	21 (42%)	13 (26%)	16 (32%)	
Pars Intermedia, Hyperplasia	3 (6%)	1 (2%)	1 (2%)	4 (8%)	
Thyroid Gland	(50)	(50)	(49)	(50)	
C-cell, Hyperplasia	34 (68%)	30 (60%)	30 (61%)	28 (56%)	
Follicular Cell, Hyperplasia	5 (10%)	1 (2%)	5 (10%)	,	
GENERAL BODY SYSTEM					
Peritoneum	(1)	(0)	(0)	(0)	
GENITAL SYSTEM					
Coagulating Gland	(0)	(0)	(0)	(1)	
Inflammation, Chronic Active	(5)	(0)	(0)	1 (100%)	
Epididymis	(50)	(50)	(50)	(50)	
Granuloma Sperm	2 (4%)	(00)	(00)	(33)	
Preputial Gland	(50)	(50)	(50)	(50)	
Inflammation, Chronic Active	3 (6%)	5 (10%)	3 (6%)	3 (6%)	
Duct, Ectasia	3 (373)	3 (1070)	0 (070)	1 (2%)	
Prostate	(50)	(50)	(50)	(50)	
Hyperplasia	12 (24%)	17 (34%)	12 (24%)	11 (22%)	
Inflammation, Chronic Active	1 (2%)	2 (4%)	4 (8%)	2 (4%)	
Seminal Vesicle	(50)	(50)	(48)	(50)	
Communicative Coloro	(00)	(30)	(10)	(00)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Inflammation, Chronic Active			2 (4%)	1 (2%)	
Testes	(50)	(50)	(50)	(50)	
Atrophy	15 (30%)	8 (16%)	5 (10%)	5 (10%)	
Edema	16 (32%)	11 (22%)	18 (36%)	11 (22%)	
Necrosis	1 (2%)				
Artery, Inflammation, Chronic Active		2 (4%)			
Interstitial Cell, Hyperplasia				1 (2%)	
HEMATOPOIETIC SYSTEM					
Bone Marrow	(50)	(50)	(49)	(50)	
Lymph Node	(12)	(5)	(7)	(5)	
Axillary, Ectasia				1 (20%)	
Axillary, Infiltration Cellular, Plasma Cell	1 (8%)				
Iliac, Ectasia	2 (17%)		1 (14%)	1 (20%)	
Iliac, Inflammation, Chronic Active			1 (14%)		
Lumbar, Ectasia	2 (17%)	2 (40%)	2 (29%)		
Lumbar, Hemorrhage	1 (8%)				
Lumbar, Infiltration Cellular, Plasma Cell	5 (42%)	1 (20%)	2 (29%)		
Pancreatic, Ectasia	1 (8%)				
Pancreatic, Inflammation, Chronic Active	1 (8%)				
Renal, Ectasia		2 (40%)		2 (40%)	
Lymph Node, Bronchial	(42)	(40)	(37)	(35)	
Ectasia				1 (3%)	
Hyperplasia, Lymphohistiocytic			10 (27%)	28 (80%)	
Lymph Node, Mandibular	(46)	(46)	(44)	(45)	
Ectasia	1 (2%)	2 (4%)			
Hyperplasia, Lymphoid		1 (2%)		1 (2%)	
Infiltration Cellular, Plasma Cell		1 (2%)	1 (2%)		
Lymph Node, Mediastinal	(46)	(45)	(50)	(49)	
Ectasia	1 (2%)				
Hyperplasia, Lymphohistiocytic			4 (8%)	29 (59%)	
Lymph Node, Mesenteric	(50)	(50)	(49)	(50)	
Ectasia	2 (4%)				
Hemorrhage				1 (2%)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Infiltration Cellular, Plasma Cell	1 (2%)				
Spleen	(50)	(50)	(50)	(50)	
Angiectasis			1 (2%)	1 (2%)	
Hematopoietic Cell Proliferation	2 (4%)		4 (8%)	2 (4%)	
Inflammation, Granulomatous		1 (2%)			
Necrosis			1 (2%)		
Thymus	(40)	(43)	(44)	(46)	
Atrophy	1 (3%)	, ,		, ,	
Cyst	, ,		1 (2%)		
INTEGUMENTARY SYSTEM					
Mammary Gland	(6)	(6)	(3)	(4)	
Galactocele	1 (17%)				
Skin	(50)	(50)	(50)	(50)	
Angiectasis	, ,	, ,		1 (2%)	
Cyst Epithelial Inclusion	1 (2%)	1 (2%)		1 (2%)	
Inflammation, Chronic Active	31 (62%)	23 (46%)	30 (60%)	22 (44%)	
Thrombosis			•	1 (2%)	
MUSCULOSKELETAL SYSTEM					
Bone	(50)	(50)	(50)	(50)	
Hyperostosis	1 (2%)				
Inflammation, Chronic Active	, ,		1 (2%)		
Skeletal Muscle	(1)	(2)	(1)	(0)	
Necrosis	`,	1 (50%)	. ,	· ,	
NERVOUS SYSTEM					
Brain	(50)	(50)	(50)	(50)	
Compression	(00)	(30)	(30)	1 (2%)	
Hemorrhage			1 (2%)	. (2/0)	
Peripheral Nerve	(5)	(4)	(6)	(6)	
i diplicial Nelve	(5)	(7)	(0)	(0)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3
Spinal Cord	(5)	(4)	(7)	(6)
RESPIRATORY SYSTEM				
Larynx	(50)	(50)	(50)	(50)
Foreign Body			1 (2%)	
Inflammation, Acute		1 (2%)		
Inflammation, Chronic				1 (2%)
Inflammation, Chronic Active		2 (4%)	2 (4%)	
Metaplasia, Squamous	1 (2%)	47 (94%)	50 (100%)	50 (100%)
Epiglottis, Fibrosis				1 (2%)
Epiglottis, Ulcer				1 (2%)
Lung	(50)	(50)	(50)	(50)
Foreign Body	1 (2%)			
Hemorrhage	1 (2%)			
Inflammation, Lymphohistiocytic	6 (12%)	14 (28%)	41 (82%)	47 (94%)
Inflammation, Acute		2 (4%)		
Metaplasia, Squamous		1 (2%)		
Alveolar Epithelium, Hyperplasia	4 (8%)	6 (12%)	11 (22%)	13 (26%)
Bronchus-associated Lymphoid Tissue, Hyperplasia, Lymphohistiocytic	·	1 (2%)	5 (10%)	19 (38%)
Nose	(50)	(50)	(50)	(50)
Foreign Body			1 (2%)	1 (2%)
Inflammation, Acute		1 (2%)	1 (2%)	1 (2%)
Inflammation, Chronic Active	4 (8%)	1 (2%)	1 (2%)	
Glands, Olfactory Epithelium, Hyperplasia	1 (2%)	39 (78%)	47 (94%)	50 (100%)
Goblet Cell, Hyperplasia		20 (40%)	25 (50%)	34 (68%)
Olfactory Epithelium, Accumulation, Hyaline Droplet	19 (38%)	50 (100%)	50 (100%)	50 (100%)
Respiratory Epithelium, Accumulation, Hyaline Droplet		17 (34%)	25 (50%)	29 (58%)
Trachea	(50)	(49)	(50)	(50)
Inflammation, Acute	. ,	1 (2%)	. ,	. ,

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
SPECIAL SENSES SYSTEM					
Ear	(0)	(0)	(1)	(0)	
Eye	(50)	(50)	(48)	(50)	
Cataract	2 (4%)	3 (6%)	1 (2%)		
Cornea, Inflammation, Chronic Active		1 (2%)		1 (2%)	
Retina, Atrophy	2 (4%)	3 (6%)	2 (4%)	1 (2%)	
Harderian Gland	(50)	(50)	(50)	(50)	
Atrophy	1 (2%)				
Hyperplasia	17 (34%)	7 (14%)	12 (24%)	6 (12%)	
Hypertrophy	2 (4%)	3 (6%)			
Zymbal's Gland	(3)	(2)	(1)	(0)	
URINARY SYSTEM					
Kidney	(50)	(50)	(49)	(50)	
Cyst	4 (8%)	1 (2%)	3 (6%)	,	
Hydronephrosis	,	1 (2%)	2 (4%)		
Infarct		,	,	1 (2%)	
Inflammation, Suppurative		3 (6%)	2 (4%)	4 (8%)	
Nephropathy	43 (86%)	38 (76%)	38 (78%)	38 (76%)	
Renal Tubule, Hyperplasia	•	, ,	• •	3 (6%)	
Transitional Epithelium, Hyperplasia	1 (2%)	1 (2%)	1 (2%)		
Urinary Bladder	(50)	(50)	(48)	(49)	
Calculus Gross Observation	, ,	1 (2%)	, ,	. ,	
Inflammation, Chronic Active		1 (2%)		1 (2%)	
Transitional Epithelium, Hyperplasia		1 (2%)		1 (2%)	

^{***} END OF MALE ***

a - Number of animals examined microscopically at site and number of animals with lesion

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

Experiment Number: 20515 - 03

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3
Disposition Summary				
Animals Initially In Study	50	50	50	50
Early Deaths				
Accidentally Killed		1		
Moribund Sacrifice	13	16	14	19
Natural Death	2			1
Survivors				
Moribund Sacrifice	1	1		1
Terminal Sacrifice	34	32	36	29
Animals Examined Microscopically	50	50	50	50
ALIMENTARY SYSTEM				
Esophagus	(50)	(50)	(50)	(50)
Inflammation, Acute	1 (2%)			
Necrosis	1 (2%)			
Intestine Large, Cecum	(49)	(50)	(50)	(49)
Intestine Large, Colon	(49)	(49)	(50)	(49)
Intestine Large, Rectum	(46)	(50)	(47)	(46)
Intestine Small, Duodenum	(49)	(50)	(50)	(49)
Muscularis, Hypertrophy, Focal			1 (2%)	
Intestine Small, Ileum	(49)	(50)	(50)	(49)
Intestine Small, Jejunum	(49)	(50)	(50)	(49)
Diverticulum	1 (2%)			
Inflammation, Acute	1 (2%)			
Inflammation, Chronic Active			1 (2%)	
Necrosis	1 (2%)			
Liver	(50)	(50)	(50)	(50)
Angiectasis	4 (8%)	5 (10%)	5 (10%)	1 (2%)
Basophilic Focus	34 (68%)	33 (66%)	35 (70%)	32 (64%)
Clear Cell Focus	14 (28%)	6 (12%)	17 (34%)	14 (28%)
Degeneration, Cystic				1 (2%)
Eosinophilic Focus	1 (2%)		1 (2%)	1 (2%)
Fatty Change	3 (6%)	1 (2%)	2 (4%)	1 (2%)
Hematopoietic Cell Proliferation	1 (2%)			

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Hepatodiaphragmatic Nodule	1 (2%)		1 (2%)		
Mixed Cell Focus	2 (4%)		` ,		
Necrosis	1 (2%)	1 (2%)	1 (2%)		
Bile Duct, Cyst	1 (2%)	1 (2%)		1 (2%)	
Bile Duct, Cyst, Multiple		4 (8%)	1 (2%)	3 (6%)	
Bile Duct, Dilatation	1 (2%)		2 (4%)	3 (6%)	
Bile Duct, Hyperplasia	5 (10%)	2 (4%)	3 (6%)	6 (12%)	
Bile Duct, Inflammation, Chronic Active	1 (2%)				
Mesentery	(7)	(8)	(8)	(5)	
Inflammation, Chronic Active				1 (20%)	
Fat, Necrosis	7 (100%)	8 (100%)	8 (100%)	4 (80%)	
Oral Mucosa	(0)	(0)	(0)	(1)	
Developmental Malformation				1 (100%)	
Inflammation, Chronic Active				1 (100%)	
Pancreas	(49)	(50)	(50)	(50)	
Atrophy	6 (12%)	4 (8%)	2 (4%)	5 (10%)	
Basophilic Focus		1 (2%)			
Salivary Glands	(50)	(50)	(50)	(50)	
Basophilic Focus	1 (2%)	1 (2%)			
Stomach, Forestomach	(50)	(50)	(50)	(50)	
Hyperplasia, Squamous	1 (2%)				
Inflammation, Acute	1 (2%)		2 (4%)		
Inflammation, Chronic Active		1 (2%)	1 (2%)		
Ulcer	1 (2%)	1 (2%)	1 (2%)	2 (4%)	
Stomach, Glandular	(50)	(50)	(50)	(50)	
Inflammation, Acute	1 (2%)				
Inflammation, Chronic Active				1 (2%)	
Mineralization	1 (2%)		1 (2%)		
Necrosis	2 (4%)		·		
Tongue	(1)	(0)	(0)	(0)	
Inflammation, Granulomatous	1 (100%)				
CARDIOVASCULAR SYSTEM					
Blood Vessel	(50)	(50)	(50)	(50)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800) CAS Number: CIMSTAR3800

Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Date Report Requested: 05/08/2013

Lab: BNW

Wistar Han RATS FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3
Heart	(50)	(50)	(50)	(50)
Cardiomyopathy	16 (32%)	17 (34%)	16 (32%)	14 (28%)
Mineralization	(=)	(5.75)	(0=70)	1 (2%)
Endocardium, Hyperplasia	1 (2%)			(=/3)
ENDOCRINE SYSTEM				
Adrenal Cortex	(50)	(50)	(50)	(50)
Accessory Adrenal Cortical Nodule	1 (2%)			
Angiectasis	1 (2%)	1 (2%)	1 (2%)	2 (4%)
Degeneration, Cystic	26 (52%)	23 (46%)	29 (58%)	35 (70%)
Hyperplasia	17 (34%)	20 (40%)	27 (54%)	22 (44%)
Hypertrophy	14 (28%)	20 (40%)	24 (48%)	21 (42%)
Metaplasia, Osseous		1 (2%)		
Necrosis	1 (2%)	1 (2%)		
Adrenal Medulla	(49)	(50)	(49)	(49)
Hyperplasia	2 (4%)	2 (4%)	1 (2%)	
Islets, Pancreatic	(49)	(50)	(50)	(50)
Hyperplasia			1 (2%)	1 (2%)
Parathyroid Gland	(47)	(44)	(40)	(42)
Pituitary Gland	(50)	(50)	(50)	(50)
Pars Distalis, Hyperplasia	8 (16%)	11 (22%)	10 (20%)	9 (18%)
Pars Intermedia, Hyperplasia	1 (2%)			1 (2%)
Thyroid Gland	(50)	(50)	(50)	(49)
C-cell, Hyperplasia	37 (74%)	44 (88%)	43 (86%)	35 (71%)
Follicular Cell, Hyperplasia	2 (4%)		1 (2%)	1 (2%)

GENERAL BODY SYSTEM

None

GENITAL SYSTEM

Clitoral Gland (50)(46)(50)(49)

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Inflammation, Chronic			1 (2%)		
Inflammation, Chronic Active	6 (12%)	4 (9%)	1 (2%)	1 (2%)	
Ovary	(50)	(50)	(50)	(50)	
Cyst	11 (22%)	9 (18%)	5 (10%)	11 (22%)	
Interstitial Cell, Hyperplasia				1 (2%)	
Uterus	(50)	(50)	(50)	(50)	
Adenomyosis			1 (2%)		
Cervix, Cyst, Squamous			1 (2%)		
Cervix, Hypertrophy	2 (4%)		1 (2%)	1 (2%)	
Endometrium, Hyperplasia, Cystic	11 (22%)	4 (8%)	9 (18%)	5 (10%)	
Vagina	(0)	(1)	(0)	(1)	
HEMATOPOIETIC SYSTEM					
Bone Marrow	(50)	(50)	(50)	(50)	
Lymph Node	(2)	(2)	(1)	(0)	
Lumbar, Ectasia		1 (50%)			
Lumbar, Hemorrhage		1 (50%)			
Lumbar, Hyperplasia, Lymphoid			1 (100%)		
Lumbar, Pigmentation, Hemosiderin	1 (50%)				
Renal, Ectasia	1 (50%)				
Renal, Pigmentation, Hemosiderin	1 (50%)				
Lymph Node, Bronchial	(38)	(35)	(32)	(35)	
Hyperplasia, Lymphohistiocytic			7 (22%)	30 (86%)	
Lymph Node, Mandibular	(49)	(47)	(43)	(49)	
Ectasia		1 (2%)		2 (4%)	
Lymph Node, Mediastinal	(49)	(46)	(45)	(47)	
Hemorrhage	` '		1 (2%)	• •	
Hyperplasia, Lymphohistiocytic			4 (9%)	23 (49%)	
Hyperplasia, Lymphoid			, ,	1 (2%)	
Lymph Node, Mesenteric	(49)	(50)	(50)	(50)	
Ectasia	1 (2%)	1 (2%)	` '	1 (2%)	
Spleen	(50)	(50)	(50)	(50)	
Hematopoietic Cell Proliferation	1 (2%)	2 (4%)	3 (6%)	1 (2%)	
Thymus	(48)	(48)	(45)	(45)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Cyst			1 (2%)		
Hyperplasia, Tubular		1 (2%)			
INTEGUMENTARY SYSTEM					
Mammary Gland	(50)	(50)	(50)	(50)	
Galactocele	3 (6%)	1 (2%)	2 (4%)	1 (2%)	
Hyperplasia	1 (2%)			1 (2%)	
Inflammation, Chronic Active				2 (4%)	
Skin	(50)	(50)	(50)	(50)	
Angiectasis	1 (2%)				
Inflammation, Chronic Active	2 (4%)	4 (8%)	5 (10%)	4 (8%)	
Subcutaneous Tissue, Hemorrhage		1 (2%)			
Subcutaneous Tissue, Necrosis		1 (2%)			
MUSCULOSKELETAL SYSTEM					
Bone	(50)	(50)	(50)	(50)	
Callus	1 (2%)	, ,	` ,	, ,	
Skeletal Muscle	(0)	(1)	(0)	(0)	
NERVOUS SYSTEM					
Brain	(50)	(50)	(50)	(50)	
Meninges, Hyperplasia	1 (2%)	(00)	(00)	1 (2%)	
Peripheral Nerve	(2)	(9)	(5)	(6)	
Spinal Cord	(2)	(9)	(5)	(6)	
	(-/		(-)		
RESPIRATORY SYSTEM					
Larynx	(50)	(50)	(50)	(50)	
Foreign Body				1 (2%)	
Inflammation, Acute				4 (8%)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Metaplasia, Squamous	1 (2%)	50 (100%)	50 (100%)	50 (100%)	
Lung	(50)	(50)	(50)	(50)	
Inflammation, Lymphohistiocytic	3 (6%)	20 (40%)	42 (84%)	50 (100%)	
Alveolar Epithelium, Hyperplasia	4 (8%)	3 (6%)	1 (2%)	7 (14%)	
Bronchus-associated Lymphoid Tissue, Hyperplasia, Lymphohistiocytic			, ,	5 (10%)	
Nose	(50)	(50)	(50)	(50)	
Inflammation, Acute				1 (2%)	
Inflammation, Chronic Active	2 (4%)		1 (2%)		
Thrombosis	1 (2%)				
Glands, Olfactory Epithelium, Hyperplasia	1 (2%)	32 (64%)	48 (96%)	49 (98%)	
Goblet Cell, Hyperplasia		25 (50%)	34 (68%)	42 (84%)	
Olfactory Epithelium, Accumulation, Hyaline Droplet	16 (32%)	50 (100%)	50 (100%)	50 (100%)	
Respiratory Epithelium, Accumulation, Hyaline Droplet	1 (2%)	24 (48%)	31 (62%)	34 (68%)	
Trachea	(49)	(50)	(50)	(50)	
SPECIAL SENSES SYSTEM					
Eye	(49)	(50)	(50)	(50)	
Cataract	(10)	(00)	1 (2%)	(55)	
Cornea, Inflammation, Chronic Active			. (= /3)	1 (2%)	
Retina, Atrophy	3 (6%)	4 (8%)	4 (8%)	2 (4%)	
Harderian Gland	(49)	(50)	(50)	(50)	
Hyperplasia	3 (6%)	3 (6%)	5 (10%)	4 (8%)	
Zymbal's Gland	(0)	(0)	(0)	(1)	
URINARY SYSTEM					
Kidney	(50)	(50)	(50)	(50)	
Cyst		1 (2%)	2 (4%)	4 (00()	
Hydronephrosis	((55()	1 (2%)		1 (2%)	
Inflammation, Suppurative	1 (2%)	2 (4%)			
Mineralization		1 (2%)		1 (2%)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: RATS/Wistar Han

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 05/08/2013 Time Report Requested: 08:51:01 First Dose M/F: 04/14/08 / 04/14/08

Wistar Han RATS FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3
Nephropathy Renal Tubule, Hyperplasia	17 (34%)	16 (32%)	19 (38%)	18 (36%) 1 (2%)
Transitional Epithelium, Hyperplasia	1 (2%)	3 (6%)	1 (2%)	. (270)
Urinary Bladder	(50)	(50)	(50)	(50)

^{***} END OF REPORT ***