Experiment Number: 20314 - 04

Species/Strain: MICE/B6C3F1/NCTR

Test Type: CHRONIC

Route: DOSED WATER

P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 12/17/2014 AVERAGE SEVERITY GRADES[b]

Glycidamide

CAS Number: 5694-00-8

Time Report Requested: 07:40:57

First Dose M/F: 06/02/05 / 06/02/05

Lab: NCTR

NTP Study Number: C20314

Lock Date: 10/03/2011

Cage Range: ALL

Date Range: ALL

Reasons For Removal: ALL

Removal Date Range: ALL

Treatment Groups: Include ALL

Study Gender: Both

TDMSE Version: 3.0.2.2_002

PWG Approval Date: NONE

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE MALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER |
|--------------------------------------|-------------|-------------|--------------|---------------|---------------|
| Disposition Summary | | | | | |
| Animals Initially In Study | 48 | 48 | 48 | 48 | 48 |
| Early Deaths | | | | | |
| Moribund Sacrifice | 11 | 16 | 11 | 4 | |
| Natural Death | 9 | 3 | 2 | 3 | 1 |
| Survivors | | | | | |
| Moribund Sacrifice | 2 | 2 | 1 | | 2 |
| Natural Death | 1 | 1 | | | |
| Terminal Sacrifice | 25 | 26 | 34 | 41 | 45 |
| Animals Examined Microscopically | 48 | 48 | 48 | 48 | 48 |
| ALIMENTARY SYSTEM | | | | | |
| Esophagus | (46) | (47) | (47) | (48) | (47) |
| Gallbladder | (39) | (45) | (46) | (45) | (46) |
| Infiltration Cellular, Lymphocyte | | 1 [3.0] | | | |
| Intestine Large, Cecum | (39) | (44) | (46) | (45) | (47) |
| Hyperplasia, Lymphoid | 2 [2.0] | 4 [1.8] | 4 [2.0] | 3 [2.0] | 1 [2.0] |
| Intestine Large, Colon | (39) | (45) | (47) | (45) | (47) |
| Intestine Large, Rectum | (39) | (45) | (47) | (45) | (47) |
| Intestine Small, Duodenum | (39) | (44) | (46) | (45) | (47) |
| Epithelium, Hyperplasia | 1 [3.0] | | | | |
| Intestine Small, Ileum | (39) | (44) | (46) | (45) | (47) |
| Hyperplasia, Lymphoid | 2 [2.0] | | | 1 [2.0] | |
| Intestine Small, Jejunum | (39) | (44) | (46) | (45) | (47) |
| Hyperplasia, Lymphoid | 1 [2.0] | | | 1 [2.0] | 1 [3.0] |
| Epithelium, Hyperplasia | | | 1 [4.0] | | |
| Liver | (48) | (47) | (47) | (47) | (47) |
| Angiectasis | 1 [1.0] | | | | |
| Basophilic Focus | 2 | 4 | 2 | | |
| Basophilic Focus, Multiple | 1 | | | | |
| Cyst | | 1 [2.0] | | | |
| Eosinophilic Focus | 2 | | | | |
| Hematopoietic Cell Proliferation | 1 [3.0] | | 2 [1.0] | 1 [1.0] | 2 [2.0] |
| Infiltration Cellular, Lymphocyte | | 3 [2.0] | 1 [1.0] | 1 [1.0] | 2 [1.0] |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE MALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|--|-------------|-------------|--------------|---------------|---------------|--|
| Infiltration Cellular, Polymorphonuclear | 1 [4.0] | | | | | |
| Inflammation, Suppurative | | | | 1 [4.0] | | |
| Inflammation, Chronic Active | 1 [1.0] | 1 [1.0] | | 3 [1.3] | 1 [1.0] | |
| Mineralization | | 1 [2.0] | | | | |
| Mixed Cell Focus | | | 1 | | | |
| Necrosis | 3 [3.7] | 3 [3.3] | 1 [3.0] | 3 [3.0] | | |
| Tension Lipidosis | | 1 [3.0] | | 1 [2.0] | | |
| Vacuolization Cytoplasmic | | 1 [1.0] | 1 [1.0] | | 2 [1.0] | |
| Bile Duct, Hyperplasia | | | | | 1 [3.0] | |
| Kupffer Cell, Hyperplasia | 1 [4.0] | | | | | |
| Oval Cell, Hyperplasia | 1 [3.0] | | | | 1 [3.0] | |
| Mesentery | (1) | (1) | (0) | (1) | (2) | |
| Hemorrhage | | | | 1 [4.0] | • • | |
| Fat, Necrosis | | 1 [3.0] | | | 1 [3.0] | |
| Oral Mucosa | (2) | (0) | (0) | (1) | (0) | |
| Pancreas | (45) | (47) | (47) | (46) | (47) | |
| Cyst | , , | 1 [4.0] | , , | , , | . , | |
| Infiltration Cellular, Lymphocyte | 3 [1.0] | 1 [1.0] | | 3 [1.3] | 7 [1.0] | |
| Acinus, Degeneration | 3 [2.0] | 6 [1.0] | 4 [1.3] | 1 [2.0] | 4 [1.5] | |
| Duct, Dilatation | 2 [3.0] | | | | • • | |
| Salivary Glands | (44) | (46) | (47) | (47) | (47) | |
| Infiltration Cellular, Lymphocyte | 18 [1.2] | 17 [1.1] | 20 [1.1] | 22 [1.3] | 26 [1.4] | |
| Stomach, Forestomach | (41) | (45) | (48) | (45) | (47) | |
| Cyst Epithelial Inclusion | , | , | 2 | , | , | |
| Ulcer | | | | 1 [2.0] | | |
| Epithelium, Hyperplasia | 12 [2.0] | 5 [2.2] | 5 [1.8] | 2 [2.0] | 5 [1.6] | |
| Serosa, Necrosis | | 1 [3.0] | | | • • | |
| Stomach, Glandular | (38) | (44) | (46) | (45) | (47) | |
| Epithelium, Hyperplasia | 1 [2.0] | , | , | 1 [3.0] | , | |
| Tongue | (0) | (1) | (0) | (0) | (0) | |
| CARDIOVASCULAR SYSTEM | | | | | | |
| Blood Vessel | (47) | (47) | (47) | (48) | (47) | |
| Heart | (46) | (47) | (47) | (48) | (47) | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE MALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|--|-------------|-------------|--------------|---------------|---------------|--|
| Infiltration Cellular, Polymorphonuclear | 1 [4.0] | | | | | |
| ENDOCRINE SYSTEM | | | | | | |
| Adrenal Cortex | (44) | (45) | (47) | (47) | (47) | |
| Accessory Adrenal Cortical Nodule | | 2 | 2 | 1 | | |
| Subcapsular, Hyperplasia | 36 [1.4] | 39 [1.2] | 45 [1.2] | 38 [1.3] | 43 [1.3] | |
| Adrenal Medulla | (43) | (45) | (47) | (46) | (47) | |
| Hyperplasia | | | | | 1 [3.0] | |
| Islets, Pancreatic | (44) | (47) | (47) | (47) | (47) | |
| Hyperplasia | 2 [2.0] | 2 [2.5] | | 2 [2.0] | 1 [2.0] | |
| Infiltration Cellular, Lymphocyte | | 1 [2.0] | | | | |
| Parathyroid Gland | (42) | (42) | (43) | (48) | (43) | |
| Cyst | 2 [1.5] | 1 [2.0] | 1 [3.0] | | 2 [2.0] | |
| Pituitary Gland | (42) | (42) | (47) | (44) | (45) | |
| Pars Distalis, Cyst | 1 [1.0] | • | 2 [1.5] | | 1 [2.0] | |
| Thyroid Gland | (43) | (44) | (47) | (48) | (47) | |
| Cyst | | | 1 [2.0] | | • | |
| Infiltration Cellular, Lymphocyte | | | - · | | 1 [1.0] | |
| Follicle, Degeneration | 4 [1.8] | 2 [1.0] | 2 [1.0] | 4 [1.3] | 2 [1.5] | |
| GENERAL BODY SYSTEM | | | | | | |
| Tissue NOS | (1) | (1) | (0) | (0) | (0) | |
| GENITAL SYSTEM | | | | | | |
| Epididymis | (43) | (46) | (47) | (47) | (47) | |
| Exfoliated Germ Cell | 1 [3.0] | (-10) | 1 [3.0] | (77) | (71) | |
| Hypospermia | 3 [2.3] | | 3 [3.0] | | 2 [2.0] | |
| Infiltration Cellular, Lymphocyte | ل [۲.۵] | | ა [ა.ს] | 1 [2.0] | ۷ [۲.۷] | |
| Infiltration Cellular, Polymorphonuclear | 1 [4.0] | | | ا رک.ن | | |
| Inflammation, Chronic | 3 [2.7] | | 2 [2.5] | 1 [2.0] | | |
| Inflammation, Chronic Active | J [Z.1] | | ک [ک.ن] | 1 [2.0] | 1 [1.0] | |
| illiammation, Chronic Active | | | | | 1 [1.0] | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE MALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|---|-------------|-------------|--------------|---------------|---------------|--|
| Spermatocele | 1 [3.0] | | | | | |
| Penis | (0) | (0) | (1) | (0) | (0) | |
| Inflammation, Chronic Active | | | 1 [2.0] | | | |
| Ulcer | | | 1 [2.0] | | | |
| Preputial Gland | (44) | (46) | (46) | (47) | (47) | |
| Cyst | 4 [2.8] | 1 [2.0] | 5 [2.4] | 2 [1.5] | 3 [2.3] | |
| Degeneration | 9 [3.1] | 12 [3.3] | 5 [3.8] | 10 [2.9] | 4 [2.8] | |
| Hyperkeratosis | | | 1 [4.0] | 1 [3.0] | 1 [2.0] | |
| Infiltration Cellular, Lymphocyte | 1 [3.0] | 4 [2.5] | 1 [2.0] | | 3 [1.0] | |
| Inflammation, Suppurative | 7 [2.7] | 3 [2.0] | 2 [2.5] | 5 [2.4] | | |
| Inflammation, Chronic Active | 2 [3.0] | | | 1 [1.0] | 1 [1.0] | |
| Duct, Dilatation | 4 [3.5] | | 1 [4.0] | | | |
| Prostate | (41) | (45) | (47) | (47) | (47) | |
| Inflammation, Suppurative | , , | | 1 [3.0] | , , | , , | |
| Inflammation, Chronic Active | 1 [4.0] | | | | | |
| Epithelium, Hyperplasia | 1 [3.0] | | | | | |
| Seminal Vesicle | (41) | (45) | (47) | (46) | (47) | |
| Atrophy | , , | 1 [3.0] | ` ' | , , | , , | |
| Lumen, Dilatation | 1 [3.0] | 1 [2.0] | 2 [1.5] | 1 [2.0] | | |
| Testes | (42) | (46) | (46) | (46) | (47) | |
| Mineralization | , , | 1 [1.0] | ` ' | , , | , , | |
| Seminiferous Tubule, Degeneration | 5 [2.4] | 2 [1.5] | 3 [3.3] | | 2 [2.0] | |
| HEMATOPOIETIC SYSTEM | | | | | | |
| Bone Marrow | (44) | (46) | (47) | (47) | (47) | |
| Hyperplasia | 8 [2.5] | 5 [2.4] | 5 [2.8] | 3 [2.0] | 3 [2.0] | |
| Lymph Node | (6) | (6) | (2) | (5) | (4) | |
| Axillary, Hyperplasia, Lymphoid | 1 [3.0] | | 1 [2.0] | 1 [2.0] | | |
| Iliac, Hyperplasia, Lymphoid | 1 [3.0] | 1 [1.0] | | | | |
| Iliac, Infiltration Cellular, Polymorphonuclear | 1 [3.0] | | | | | |
| Inguinal, Hyperplasia, Lymphoid | 1 [2.0] | 2 [1.5] | 1 [2.0] | | | |
| Inguinal, Infiltration Cellular, Plasma Cell | 1 [2.0] | | | | | |
| Lumbar, Hyperplasia, Lymphoid | | | 1 [1.0] | | 1 [4.0] | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| 57BL/6N XC3H/HEN MTV-NCTR MICE MALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER |
|---|-------------|----------------------|----------------------|-----------------------|---------------|
| Lumbar, Infiltration Cellular, | 1 [4.0] | | | | |
| Polymorphonuclear | | 4 [0.0] | | | 4 [4 0] |
| Mediastinal, Hyperplasia, Lymphoid | 4 [0 0] | 1 [3.0] | | | 1 [4.0] |
| Renal, Hyperplasia, Lymphoid | 1 [3.0] | | | | 1 [4.0] |
| Renal, Infiltration Cellular, Polymorphonuclear | 1 [4.0] | (45) | (47) | (47) | (4=) |
| Lymph Node, Mandibular | (43) | (45) | (47) | (47) | (47) |
| Hyperplasia, Lymphoid | 7 [2.0] | 10 [2.4] | 5 [1.8] | 4 [1.3] | 7 [2.0] |
| Infiltration Cellular, Histiocyte | 2 [2.0] | 1 [2.0] | | | |
| Infiltration Cellular, Mast Cell | 2 [3.0] | 1 [2.0] | 1 [2.0] | | |
| Infiltration Cellular, Plasma Cell | 4 [2.3] | 3 [2.0] | 1 [2.0] | | 1 [3.0] |
| Infiltration Cellular, Polymorphonuclear | 1 [4.0] | | | | |
| Pigmentation | 1 [1.0] | | 1 [1.0] | | |
| Sinus, Dilatation | 1 [3.0] | | | | |
| Lymph Node, Mesenteric | (42) | (45) | (47) | (46) | (47) |
| Angiectasis | 3 [2.0] | 5 [3.2] | 3 [2.0] | 5 [2.2] | 9 [2.4] |
| Hematopoietic Cell Proliferation | 4 [2.5] | | | | |
| Hemorrhage | 9 [2.8] | 4 [2.3] | 4 [2.0] | 12 [2.2] | 5 [2.2] |
| Hyperplasia, Lymphoid | 12 [2.5] | 16 [1.8] | 21 [2.3] | 12 [2.1] | 25 [2.1] |
| Infiltration Cellular, Histiocyte | 5 [3.0] | 5 [3.0] | 3 [3.0] | 4 [3.3] | 5 [2.8] |
| Infiltration Cellular, Mast Cell | 2 [2.0] | | 1 [2.0] | 2 [2.0] | |
| Infiltration Cellular, Polymorphonuclear | 1 [4.0] | | | | 1 [2.0] |
| Sinus, Dilatation | 1 [4.0] | | | 1 [2.0] | 1 |
| Spleen | (44) | (46) | (47) | (47) | (47) |
| Congestion | 1 [4.0] | (/ | (· · / | (, | \···/ |
| Depletion Lymphoid | , [,,•] | 2 [2.5] | | | |
| Hematopoietic Cell Proliferation | 17 [3.1] | 14 [3.1] | 12 [3.3] | 6 [3.2] | 6 [3.7] |
| Hyperplasia, Lymphoid | 23 [2.6] | 22 [2.5] | 22 [2.4] | 31 [2.4] | 34 [2.7] |
| Inflammation, Granulomatous | 23 [2.0] | کد [د.ن] | 22 [2.4] 1 [4.0] | 31 [2. 4] | J4 [Z./] |
| Necrosis | | 1 [4.0] | ı [4 .0] | | |
| Pigmentation | 1 [2.0] | ı [4 .0] | | | |
| • | | (20) | (42) | (45) | (42) |
| Thymus | (34) | (39) | (43) | (45) | (43) |
| Atrophy | 16 [3.1] | 21 [3.1] | 17 [3.2] | 25 [2.7] | 28 [2.7] |
| Hyperplasia, Lymphoid | | 1 [2.0] | 1 [2.0] | | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57
First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE MALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|--------------------------------------|-------------|-------------|--------------|---------------|---------------|--|
| INTEGUMENTARY SYSTEM | | | | | | |
| Mammary Gland | (1) | (2) | (0) | (1) | (3) | |
| Skin | (46) | (47) | (47) | (48) | (47) | |
| Fibrosis | | 1 [2.0] | | 1 [3.0] | | |
| Inflammation, Chronic Active | 1 [2.0] | 1 [2.0] | 1 [2.0] | | | |
| Metaplasia, Osseous | | | | | 1 [4.0] | |
| Mineralization | | 1 [4.0] | | | | |
| Ulcer | 1 [2.0] | | | | | |
| Epithelium, Hyperplasia | 1 [2.0] | | 1 [2.0] | | | |
| Sebaceous Gland, Hyperplasia | | 1 [2.0] | · - | | | |
| MUSCULOSKELETAL SYSTEM | | | | | | |
| Bone, Femur | (47) | (48) | (48) | (48) | (48) | |
| Skeletal Muscle | (47) | (47) | (47) | (47) | (48) | |
| Degeneration | | 1 [3.0] | | | | |
| NERVOUS SYSTEM | | | | | | |
| Brain, Brain Stem | (42) | (46) | (46) | (47) | (47) | |
| Brain, Cerebellum | (42) | (46) | (46) | (47) | (47) | |
| Hemorrhage | , , | , , | 1 [1.0] | ` ' | , | |
| Brain, Cerebrum | (42) | (46) | (46) | (47) | (47) | |
| Cyst | , , | , , | , , | 1 [1.0] | , | |
| Mineralization | 20 [1.2] | 29 [1.2] | 32 [1.2] | 30 [1.2] | 33 [1.2] | |
| Meninges, Arteritis | | | | • • | 1 [1.0] | |
| Peripheral Nerve, Sciatic | (42) | (46) | (47) | (47) | (47) | |
| Axon, Degeneration | 32 [1.3] | 30 [1.3] | 30 [1.3] | 37 [1.1] | 36 [1.1] | |
| Spinal Cord, Cervical | (39) | (44) | (46) | (46) | (47) | |
| Axon, Degeneration | 12 [1.0] | 11 [1.0] | 3 [1.0] | 8 [1.0] | 8 [1.0] | |
| Nerve, Degeneration | | | 1 [1.0] | | | |
| Spinal Cord, Lumbar | (39) | (45) | (46) | (46) | (47) | |
| Axon, Degeneration | 1 [1.0] | , , | 4 [1.0] | 10 [1.0] | 9 [1.0] | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 **First Dose M/F:** 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE MALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER |
|---|-------------|-------------|--------------|---------------|---------------|
| Nerve, Degeneration | 34 [1.6] | 35 [1.3] | 37 [1.2] | 41 [1.5] | 39 [1.2] |
| Spinal Cord, Thoracic | (39) | (45) | (46) | (46) | (47) |
| Axon, Degeneration | 26 [1.0] | 24 [1.0] | 23 [1.0] | 28 [1.0] | 37 [1.0] |
| Nerve, Degeneration | | | 1 [1.0] | 1 [1.0] | 1 [1.0] |
| RESPIRATORY SYSTEM | | | | , | , |
| Lung | (47) | (47) | (47) | (46) | (47) |
| Congestion | | | 1 [4.0] | | |
| Hemorrhage | 1 [2.0] | | | | |
| Infiltration Cellular, Histiocyte | 4 [3.8] | 1 [3.0] | | 1 [1.0] | |
| Infiltration Cellular, Lymphocyte | | | 1 [1.0] | | |
| Inflammation, Chronic | 1 [1.0] | | | | |
| Alveolar Epithelium, Hyperplasia | 6 [2.5] | 3 [3.3] | 4 [2.3] | 1 [2.0] | |
| Alveolus, Infiltration Cellular, Histiocyte | 1 [1.0] | | | | |
| Nose | (45) | (47) | (47) | (46) | (47) |
| Amyloid Deposition | | | 1 [2.0] | | |
| Crystals | 1 [1.0] | 1 [3.0] | 2 [1.5] | | |
| Foreign Body | | 1 | | | |
| Hyaline Droplet | 1 [1.0] | 2 [1.5] | 11 [1.6] | 6 [2.0] | 13 [1.8] |
| Inflammation, Suppurative | 1 [1.0] | 1 [2.0] | | 1 [2.0] | |
| Trachea | (45) | (46) | (46) | (48) | (47) |
| SPECIAL SENSES SYSTEM | | | | | |
| Eye | (42) | (44) | (46) | (45) | (47) |
| Cataract | 17 [2.0] | 8 [2.3] | 7 [2.1] | 3 [1.0] | 1 [4.0] |
| Cornea, Inflammation, Chronic | 1 [1.0] | | | | |
| Cornea, Inflammation, Chronic Active | 7 [2.3] | | 2 [2.5] | | |
| Cornea, Ulcer | 3 [3.3] | | 1 [4.0] | | |
| Harderian Gland | (47) | (46) | (47) | (47) | (47) |
| Cyst | 1 [4.0] | | | | |
| Infiltration Cellular, Lymphocyte | | | 1 [1.0] | | 4 [1.3] |
| Epithelium, Hyperplasia | 1 [2.0] | 2 [2.0] | 2 [2.5] | 1 [3.0] | |
| Zymbal's Gland | (1) | (0) | (0) | (0) | (0) |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Experiment Number: 20314 - 04

P18: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE (a) WITH Date Report Requested: 12/17/2014 AVERAGE SEVERITY GRADES[b]

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57
First Dose M/F: 06/02/05 / 06/02/05

Lab: NCTR

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

| 57BL/6N XC3H/HEN MTV-NCTR MICE MALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER |
|--|-------------|-------------|--------------|---------------|---------------|
| Keratin Cyst | 1 | | | | |
| RINARY SYSTEM | | | | | |
| Kidney | (46) | (46) | (47) | (47) | (47) |
| Autolysis | 1 [4.0] | 1 [4.0] | | | |
| Cyst | | | | | 1 [2.0] |
| Degeneration | 1 [4.0] | | | | |
| Hyaline Droplet | 2 [3.5] | 3 [3.7] | 1 [3.0] | | |
| Hydronephrosis | 1 [4.0] | | | | 1 [2.0] |
| Infarct | 1 [2.0] | 1 [2.0] | | | 1 [2.0] |
| Infiltration Cellular, Lymphocyte | 9 [1.4] | 8 [1.5] | 19 [1.1] | 12 [1.1] | 20 [1.5] |
| Infiltration Cellular, Polymorphonuclear | 1 [4.0] | | | | |
| Inflammation, Chronic | | | | | 1 [3.0] |
| Nephropathy | 16 [1.1] | 12 [1.1] | 14 [1.3] | 18 [1.1] | 15 [1.0] |
| Urinary Bladder | (41) | (45) | (47) | (46) | (47) |
| Infiltration Cellular, Lymphocyte | 1 [1.0] | 3 [1.0] | 3 [1.0] | 4 [1.0] | 7 [1.0] |
| Infiltration Cellular, Polymorphonuclear | 1 [4.0] | | | | |
| Inflammation, Suppurative | 1 [4.0] | | | | |
| Lumen, Dilatation | 2 [3.5] | 1 [3.0] | 3 [4.0] | 2 [4.0] | 3 [3.7] |

^{***} END OF MALE ***

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE FEMALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER |
|--|-------------|-------------|--------------|---------------|---------------|
| Disposition Summary | | | | | |
| Animals Initially In Study | 48 | 48 | 48 | 48 | 48 |
| Early Deaths | | | | | |
| Moribund Sacrifice | 32 | 7 | 6 | 2 | 2 |
| Natural Death | 6 | 4 | 2 | 2 | 1 |
| Survivors | | | | | |
| Moribund Sacrifice | 2 | 5 | 2 | | 2 |
| Natural Death | | 1 | | 2 | 2 |
| Terminal Sacrifice | 8 | 31 | 38 | 42 | 41 |
| Animals Examined Microscopically | 48 | 48 | 48 | 48 | 47 |
| ALIMENTARY SYSTEM | | | | | |
| Esophagus | (45) | (47) | (46) | (48) | (46) |
| Gallbladder | (41) | (44) | (47) | (44) | (45) |
| Infiltration Cellular, Lymphocyte | | | | 1 [1.0] | |
| Intestine Large, Cecum | (42) | (45) | (46) | (44) | (45) |
| Hyperplasia, Lymphoid | 1 [2.0] | 4 [2.0] | | | 2 [2.5] |
| Intestine Large, Colon | (43) | (45) | (47) | (45) | (45) |
| Intestine Large, Rectum | (43) | (45) | (46) | (45) | (45) |
| Intestine Small, Duodenum | (42) | (45) | (46) | (44) | (45) |
| Hyperplasia, Lymphoid | 1 [3.0] | | | | |
| Intestine Small, Ileum | (42) | (45) | (46) | (44) | (45) |
| Hyperplasia, Lymphoid | | 1 [2.0] | | | |
| Inflammation, Suppurative | | | | 1 [4.0] | |
| Ulcer | | | | 1 [4.0] | |
| Intestine Small, Jejunum | (42) | (45) | (46) | (44) | (45) |
| Hyperplasia, Lymphoid | | 1 [1.0] | 1 [3.0] | 1 [3.0] | |
| Liver | (43) | (46) | (47) | (48) | (47) |
| Angiectasis | 5 [2.4] | · | 1 [2.0] | · | |
| Basophilic Focus | 4 | 1 | 3 | | 2 |
| Cyst | 1 [2.0] | 1 [4.0] | | | 3 [2.7] |
| Eosinophilic Focus | | - | | 2 | |
| Eosinophilic Focus, Multiple | 1 | | | | |
| Hematopoietic Cell Proliferation | 6 [1.8] | 5 [1.4] | 3 [1.0] | 11 [1.5] | 6 [1.0] |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE FEMALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|---|-------------|-------------|--------------|---------------|---------------|--|
| Hemorrhage | | | | | 1 [2.0] | |
| Infiltration Cellular, Lymphocyte | 2 [2.0] | 4 [1.5] | 3 [1.3] | 3 [1.3] | 4 [1.0] | |
| Inflammation, Chronic Active | 1 [1.0] | 5 [1.0] | | | 1 [4.0] | |
| Necrosis | 5 [2.2] | | | | | |
| Tension Lipidosis | 1 [2.0] | | | | | |
| Vacuolization Cytoplasmic | 3 [2.0] | 3 [1.7] | 1 [2.0] | 1 [2.0] | 1 [2.0] | |
| Centrilobular, Degeneration | 1 [3.0] | | | | | |
| Pancreas | (44) | (46) | (47) | (48) | (45) | |
| Infiltration Cellular, Lymphocyte | 3 [1.0] | 9 [1.1] | 10 [1.2] | 10 [1.1] | 9 [1.2] | |
| Necrosis | 2 [4.0] | | | | | |
| Acinus, Degeneration | 2 [1.5] | | 3 [2.3] | 6 [1.7] | 2 [3.0] | |
| Duct, Dilatation | | | | 2 [4.0] | | |
| Salivary Glands | (44) | (47) | (47) | (47) | (45) | |
| Infiltration Cellular, Lymphocyte | 8 [1.4] | 28 [1.3] | 28 [1.3] | 30 [1.2] | 29 [1.4] | |
| Stomach, Forestomach | (44) | (45) | (47) | (45) | (45) | |
| Cyst Epithelial Inclusion | | | | 1 | | |
| Ulcer | 1 [2.0] | 1 [2.0] | 1 [2.0] | | 1 [2.0] | |
| Epithelium, Hyperplasia | 5 [2.4] | 11 [2.5] | 10 [2.4] | 4 [2.8] | 4 [3.0] | |
| Stomach, Glandular | (43) | (45) | (47) | (45) | (45) | |
| Epithelium, Hyperplasia | | 1 [2.0] | | | | |
| CARDIOVASCULAR SYSTEM | | | | | | |
| Blood Vessel | (45) | (47) | (47) | (48) | (46) | |
| Heart | (45) | (47) | (47) | (48) | (46) | |
| Cardiomyopathy | 1 [1.0] | | | 1 [1.0] | , , | |
| Inflammation, Chronic | | | | 1 [1.0] | | |
| Pericardium, Inflammation, Chronic Active | 1 [2.0] | | | | | |
| ENDOCRINE SYSTEM | | | | | | |
| Adrenal Cortex | (44) | (47) | (47) | (48) | (45) | |
| Accessory Adrenal Cortical Nodule | | 2 | | | | |
| Angiectasis | | 1 [2.0] | | | | |
| Cyst | 1 [4.0] | | | | | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 **First Dose M/F:** 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE FEMALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER |
|--|-------------|-------------|--------------|---------------|---------------|
| Hypertrophy | | | 1 [3.0] | 1 [2.0] | |
| Subcapsular, Hyperplasia | 43 [2.2] | 46 [2.1] | 47 [2.2] | 48 [2.1] | 45 [2.1] |
| Adrenal Medulla | (43) | (47) | (47) | (45) | (45) |
| Islets, Pancreatic | (43) | (46) | (47) | (48) | (45) |
| Hyperplasia | | 2 [2.0] | 3 [1.7] | 2 [2.0] | |
| Parathyroid Gland | (39) | (40) | (46) | (45) | (41) |
| Cyst | 2 [1.5] | 2 [2.0] | 2 [2.0] | | 4 [2.3] |
| Hyperplasia | 1 [2.0] | | | | |
| Inflammation, Chronic Active | | | 1 [1.0] | | |
| Pituitary Gland | (43) | (47) | (43) | (45) | (45) |
| Pars Distalis, Cyst | | | 1 [4.0] | 1 [4.0] | |
| Pars Distalis, Hyperplasia | 3 [1.7] | 1 [2.0] | 2 [2.0] | | 3 [2.3] |
| Thyroid Gland | (44) | (46) | (46) | (47) | (45) |
| Cyst | ` , | | , , | 1 [2.0] | 1 [3.0] |
| Ectopic Thymus | 1 [2.0] | 1 [1.0] | 3 [1.3] | 2 [2.0] | 1 [1.0] |
| Infiltration Cellular, Lymphocyte | | | | 3 [1.0] | 1 [2.0] |
| Inflammation, Chronic Active | | 1 [1.0] | | 1 [1.0] | |
| Follicle, Degeneration | 3 [1.7] | 6 [1.2] | 7 [1.0] | 5 [1.4] | 3 [1.3] |
| GENERAL BODY SYSTEM | | | | | |
| Tissue NOS | (1) | (0) | (0) | (0) | (0) |
| GENITAL SYSTEM | | | | | |
| Clitoral Gland | (41) | (47) | (47) | (48) | (44) |
| Degeneration | 36 [3.9] | 47 [4.0] | 47 [3.9] | 48 [3.9] | 43 [4.0] |
| Inflammation, Suppurative | | • • | • • | | 1 [2.0] |
| Inflammation, Chronic Active | 1 [2.0] | | | | 1 [2.0] |
| Ovary | (44) | (46) | (47) | (47) | (45) |
| Angiectasis | 1 [4.0] | 4 [2.8] | , , | , , | 1 [4.0] |
| Atrophy | 31 [3.4] | 39 [3.8] | 46 [3.9] | 42 [3.9] | 44 [4.0] |
| Cyst | 15 [2.9] | 21 [3.0] | 23 [2.8] | 16 [2.6] | 14 [2.6] |
| Hematocyst | 1 [4.0] | | | | |
| Hemorrhage | 3 [3.7] | 2 [2.5] | 3 [2.7] | 1 [2.0] | 1 [4.0] |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 **First Dose M/F:** 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE FEMALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER |
|--|-------------|-------------|--------------|---------------|---------------|
| Hyperplasia, Tubulostromal | 1 [3.0] | | | | |
| Inflammation, Chronic | | | | | 1 [4.0] |
| Mineralization | 1 [3.0] | | | | 1 [4.0] |
| Thrombosis | 1 [4.0] | 1 [4.0] | 1 [3.0] | | 1 [4.0] |
| Bilateral, Cyst | 3 [3.0] | 1 [2.0] | 2 [4.0] | 1 [4.0] | |
| Uterus | (45) | (47) | (47) | (48) | (45) |
| Angiectasis | 2 [2.5] | 2 [3.5] | | 2 [4.0] | 1 [3.0] |
| Fibrosis | | | | 1 [4.0] | |
| Hemorrhage | 1 [4.0] | 1 [2.0] | | | |
| Hydrometra | | | | 1 [4.0] | 1 [4.0] |
| Inflammation, Suppurative | | | 1 [2.0] | | |
| Necrosis | 1 [4.0] | | | | |
| Thrombosis | 2 [2.5] | | | 1 [4.0] | |
| Endometrium, Hyperplasia, Cystic | 36 [2.3] | 45 [2.3] | 47 [2.6] | 45 [2.8] | 44 [2.9] |
| Endometrium, Hyperplasia, Glandular, Focal | | | 1 [3.0] | | |
| Vagina | (1) | (0) | (0) | (0) | (0) |
| HEMATOPOIETIC SYSTEM | | | | | |
| Bone Marrow | (43) | (45) | (47) | (47) | (46) |
| Hyperplasia | 20 [2.7] | 6 [2.8] | 5 [2.2] | 2 [2.5] | , |
| Lymph Node | (12) | (8) | (5) | (2) | (8) |
| Lumbar, Hyperplasia, Lymphoid | 2 [2.5] | 1 [2.0] | () | () | 1 [2.0] |
| Lumbar, Infiltration Cellular, Histiocyte | 1 [2.0] | | | | |
| Lumbar, Infiltration Cellular, Plasma Cell | 1 [2.0] | | | | |
| Lumbar, Infiltration Cellular, Polymorphonuclear | 1 [3.0] | 1 [2.0] | | | |
| Mediastinal, Angiectasis | 1 [3.0] | | | | |
| Mediastinal, Hyperplasia, Lymphoid | | | 2 [3.0] | | 1 [3.0] |
| Pancreatic, Hyperplasia, Lymphoid | | 1 [2.0] | r1 | | 11 |
| Renal, Hyperplasia, Lymphoid | 1 [3.0] | 1 | | | |
| Renal, Infiltration Cellular, Plasma Cell | 1 [3.0] | | | | |
| Renal, Infiltration Cellular, Polymorphonuclear | 1 [3.0] | | | 1 [2.0] | |
| Lymph Node, Mandibular | (43) | (47) | (47) | (48) | (46) |
| Erythrophagocytosis | 1 [4.0] | () | (/ | () | (. •) |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 **First Dose M/F:** 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE FEMALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|--|-------------|-------------|--------------|---------------|---------------|--|
| Hemorrhage | | | | | 1 [2.0] | |
| Hyperplasia, Lymphoid | | 8 [2.1] | 8 [1.6] | 10 [1.8] | 13 [1.6] | |
| Infiltration Cellular, Plasma Cell | 1 [3.0] | | 4 [2.0] | 1 [3.0] | 3 [2.0] | |
| Infiltration Cellular, Polymorphonuclear | 1 [3.0] | | | 2 [2.5] | 1 [2.0] | |
| Sinus, Dilatation | 1 [3.0] | | | | | |
| Lymph Node, Mesenteric | (43) | (47) | (46) | (46) | (43) | |
| Angiectasis | | 2 [3.5] | | 2 [1.5] | | |
| Hemorrhage | 1 [2.0] | | | | | |
| Hyperplasia, Lymphoid | 5 [2.2] | 14 [2.2] | 9 [2.0] | 7 [1.4] | 14 [1.6] | |
| Infiltration Cellular, Histiocyte | | 2 [2.5] | 1 [3.0] | | 1 [3.0] | |
| Infiltration Cellular, Mast Cell | | | 1 [3.0] | | | |
| Infiltration Cellular, Plasma Cell | | | • • | 1 [2.0] | 2 [3.0] | |
| Infiltration Cellular, Polymorphonuclear | | | | 1 [2.0] | | |
| Thrombosis | 1 [2.0] | | | | | |
| Sinus, Dilatation | | | | | 1 [2.0] | |
| Spleen | (45) | (47) | (47) | (47) | (46) | |
| Accessory Spleen | () | , | 1 | , | , | |
| Depletion Lymphoid | 1 [3.0] | | | | | |
| Hematopoietic Cell Proliferation | 29 [3.4] | 14 [3.6] | 11 [3.5] | 10 [3.3] | 6 [3.0] | |
| Hemorrhage | - [-] | 11 | 1 [4.0] | - [] | - 11 | |
| Hyperplasia, Lymphoid | 12 [2.7] | 29 [2.9] | 35 [2.8] | 33 [3.0] | 35 [3.1] | |
| Necrosis | | - 1 - 1 | | | 1 [4.0] | |
| Pigmentation | | 1 [2.0] | 1 [2.0] | 3 [2.3] | 1 [3.0] | |
| Thymus | (37) | (43) | (45) | (44) | (43) | |
| Angiectasis | (- / | (- / | (- / | () | 1 [2.0] | |
| Atrophy | 18 [3.7] | 13 [2.8] | 15 [2.7] | 11 [2.5] | 16 [2.9] | |
| Hyperplasia, Lymphoid | 1 [2.0] | 5 [2.0] | 9 [2.0] | 5 [1.8] | 7 [2.1] | |
| Mineralization | | | 1 [4.0] | 1 [3.0] | | |
| INTEGUMENTARY SYSTEM | | | | | | |
| Mammary Gland | (45) | (47) | (47) | (48) | (45) | |
| Alveolus, Hyperplasia | | 2 [2.5] | | | 1 [2.0] | |
| Skin | (45) | (47) | (47) | (48) | (45) | |
| Fibrosis | | 2 [3.5] | | | | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Experiment Number: 20314 - 04

Species/Strain: MICE/B6C3F1/NCTR

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE FEMALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|--|-------------|-------------|--------------|---------------|---------------|--|
| Inflammation, Chronic Active | | 1 [4.0] | | | | |
| MUSCULOSKELETAL SYSTEM | | , | | | | |
| Bone | (1) | (2) | (1) | (0) | (0) | |
| Joint, Ligament, Degeneration | | | 1 [2.0] | | | |
| Bone, Femur | (48) | (48) | (48) | (48) | (46) | |
| Callus | 1 [3.0] | | | | | |
| Fibro-Osseous Lesion | | 1 [3.0] | | | 1 [3.0] | |
| Skeletal Muscle | (45) | (47) | (47) | (48) | (46) | |
| Atrophy | 1 [2.0] | | | | | |
| NERVOUS SYSTEM | | | | | | |
| Brain, Brain Stem | (44) | (45) | (47) | (47) | (45) | |
| Brain, Cerebellum | (44) | (45) | (47) | (47) | (45) | |
| Hemorrhage | 1 [2.0] | | | | | |
| Infiltration Cellular, Lymphocyte | 1 [1.0] | | | 1 [1.0] | | |
| Brain, Cerebrum | (44) | (45) | (47) | (47) | (45) | |
| Compression | | 1 [1.0] | | | | |
| Hemorrhage | 1 [2.0] | | | | | |
| Infarct, Focal, Chronic | 1 [1.0] | | | | | |
| Infiltration Cellular, Lymphocyte | | | | 1 [1.0] | 3 [1.0] | |
| Mineralization | 22 [1.0] | 25 [1.0] | 28 [1.0] | 36 [1.1] | 31 [1.0] | |
| Hippocampus, Neuron, Depletion | | | 1 [2.0] | | | |
| Peripheral Nerve, Sciatic | (44) | (45) | (47) | (47) | (45) | |
| Inflammation, Acute | | | 1 [2.0] | | | |
| Axon, Degeneration | 28 [1.1] | 34 [1.1] | 38 [1.1] | 38 [1.2] | 39 [1.2] | |
| Spinal Cord, Cervical | (43) | (45) | (47) | (44) | (45) | |
| Infiltration Cellular, Lymphocyte | | 1 [1.0] | 1 [1.0] | | 3 [1.0] | |
| Axon, Degeneration | 10 [1.0] | 9 [1.0] | 10 [1.0] | 9 [1.0] | 4 [1.0] | |
| Nerve, Degeneration | 1 [1.0] | | | | | |
| Spinal Cord, Lumbar | (43) | (45) | (47) | (45) | (45) | |
| Cyst | | | 1 [3.0] | | | |
| Infiltration Cellular, Lymphocyte | 1 [1.0] | 1 [1.0] | 1 [1.0] | 1 [1.0] | 6 [1.0] | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE FEMALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|--|-------------|-------------|--------------|---------------|---------------|--|
| Axon, Degeneration | 4 [1.0] | 15 [1.0] | 13 [1.0] | 13 [1.0] | 12 [1.0] | |
| Nerve, Degeneration | 24 [1.3] | 40 [1.6] | 40 [1.4] | 43 [1.2] | 43 [1.2] | |
| Spinal Cord, Thoracic | (43) | (45) | (47) | (44) | (45) | |
| Infiltration Cellular, Lymphocyte | | 1 [1.0] | 1 [1.0] | | 3 [1.0] | |
| Axon, Degeneration | 23 [1.0] | 34 [1.0] | 34 [1.0] | 32 [1.0] | 35 [1.0] | |
| Nerve, Degeneration | | 1 [1.0] | | 1 [1.0] | | |
| RESPIRATORY SYSTEM | | | | | | |
| Lung | (44) | (47) | (47) | (48) | (46) | |
| Congestion | | | | 1 [3.0] | | |
| Hemorrhage | 1 [2.0] | | | | 2 [2.5] | |
| Infiltration Cellular, Histiocyte | 3 [3.0] | 2 [1.5] | 1 [1.0] | | | |
| Infiltration Cellular, Lymphocyte | 1 [2.0] | 1 [2.0] | 2 [1.0] | 3 [1.7] | 1 [1.0] | |
| Inflammation, Chronic Active | | | 1 [1.0] | 2 [1.5] | 3 [3.0] | |
| Alveolar Epithelium, Hyperplasia | 3 [2.3] | 4 [2.0] | 2 [2.5] | 1 [2.0] | | |
| Nose | (45) | (46) | (47) | (46) | (46) | |
| Amyloid Deposition | | | 1 [1.0] | | | |
| Crystals | 1 [2.0] | | 2 [2.5] | | | |
| Hyaline Droplet | | 4 [1.5] | 15 [2.0] | 3 [1.3] | 9 [1.4] | |
| Inflammation, Suppurative | 1 [2.0] | | | 1 [2.0] | | |
| Trachea | (43) | (46) | (46) | (47) | (45) | |
| Atrophy | . , | 1 [4.0] | , , | , , | , , | |
| SPECIAL SENSES SYSTEM | | | | | | |
| Eye | (43) | (44) | (47) | (44) | (45) | |
| Cataract | 9 [1.6] | 8 [2.5] | 8 [3.1] | 2 [3.0] | 1 [1.0] | |
| Inflammation, Chronic Active | | - | - | 1 [1.0] | | |
| Phthisis Bulbi | 2 | 1 | | 1 | | |
| Cornea, Fibrosis | 1 [2.0] | | | | | |
| Cornea, Inflammation, Chronic Active | 5 [2.2] | 3 [1.3] | 1 [1.0] | 2 [2.0] | | |
| Cornea, Ulcer | 1 [3.0] | - · | - · | | | |
| Harderian Gland | (46) | (46) | (47) | (47) | (45) | |
| Degeneration, Cystic | . , | 1 [4.0] | | . , | | |

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)

Test Type: CHRONIC Route: DOSED WATER

Species/Strain: MICE/B6C3F1/NCTR

Experiment Number: 20314 - 04

Glycidamide CAS Number: 5694-00-8

Time Report Requested: 07:40:57 First Dose M/F: 06/02/05 / 06/02/05

| C57BL/6N XC3H/HEN MTV-NCTR MICE FEMALE | 0.70 GLYCID | 0.35 GLYCID | 0.175 GLYCID | 0.0875 GLYCID | CONTROL WATER | |
|--|-------------|-------------|--------------|---------------|---------------|--|
| Fibrosis | | 1 [4.0] | 1 [4.0] | | | |
| Infiltration Cellular, Lymphocyte | | 2 [1.5] | 1 [1.0] | 5 [1.2] | 3 [1.3] | |
| Epithelium, Hyperplasia | | 1 [2.0] | | 1 [3.0] | | |
| URINARY SYSTEM | | | | | | |
| Kidney | (43) | (46) | (47) | (47) | (46) | |
| Cyst | 1 [4.0] | | | | | |
| Hyaline Droplet | 3 [2.7] | | 1 [2.0] | 1 [4.0] | | |
| Hydronephrosis | 1 [4.0] | | 1 [4.0] | | | |
| Infarct | | | 1 [3.0] | | | |
| Infiltration Cellular, Lymphocyte | 13 [1.5] | 19 [1.2] | 28 [1.2] | 26 [1.3] | 23 [1.5] | |
| Inflammation, Chronic | | | 1 [4.0] | | | |
| Metaplasia, Osseous | | | | 1 [1.0] | 2 [1.5] | |
| Nephropathy | 5 [1.6] | 5 [1.0] | 3 [1.3] | 2 [1.5] | 2 [1.5] | |
| Glomerulus, Amyloid Deposition | | 1 [3.0] | | | 1 [3.0] | |
| Urinary Bladder | (42) | (46) | (47) | (46) | (45) | |
| Infiltration Cellular, Lymphocyte | 18 [1.1] | 22 [1.1] | 34 [1.1] | 28 [1.1] | 31 [1.3] | |
| Lumen, Dilatation | 1 [3.0] | 2 [3.5] | | | | |

^{***} END OF REPORT ***

a - Number of animals examined microscopically at site and number of animals with lesion b-Average severity grade(1-minimal;2-mild;3-moderate;4-marked)