

Table 7
Incidence of Cardiac Lesions at PND90.

AGE	Exposure Duration	Sex/Pathology	0	BPA 2.5	BPA 25	BPA 250	BPA 2500	BPA 25000	EE2 0.05	EE2 0.5
PND 90	Stop	Male								
		Cardiomyopathy	10/10 (1.9)	10/10 (1.5)	8/8 (1.5)	9/10 (1.3)	9/9 (2.0)	10/10 (1.6)	9/10 (1.4)	10/10 (1.6)
		Focal Fibrosis	2/10	3/10	4/8	2/10	5/9	3/10	4/10	3/10
		Inflammation/Necrosis	4/10	1/10	3/8	2/10	3/9	3/10	2/10	2/10
	Diffuse Degeration	–	–	–	2/10	–	1/10	2/10	1/10	
	Stop	Female								
		Cardiomyopathy	10/10 (1.4)	8/8 (1.4)	10/10 (1.1)	10/10 (1.1)	7/8 (0.9)	10/10 (1.5)	10/10 (1.3)	10/10 (2.0)
		Focal Fibrosis	4/10	2/8	6/10	3/10	6/8	5/10	5/10	7/10
		Inflammation/Necrosis	1/10	3/8	2/10	2/10	1/10	1/10	0/10	1/10
	Diffuse Degeration	–	–	–	1/10	–	–	–	–	
	Continuous	Male								
		Cardiomyopathy	10/10 (1.3)	9/9 (1.8)	8/8 (1.9)	10/10 (1.9)	8/8 (1.6)	9/10 (1.6)	10/10 (1.5)	10/10 (1.7)
		Focal Fibrosis	5/10	2/9	4/8	4/10	2/8	3/10	1/10	4/10
		Inflammation/Necrosis	4/10	3/9	4/8	6/10	3/8	4/10	3/10	3/10
	Diffuse Degeration	–	–	1/8	2/10	–	1/10	–	2/10	
	Continuous	Female								
Cardiomyopathy		10/10 (1.5)	9/10 (1.6)	9/10 (1.2)	10/10 (1.7)	9/9 (1.4)	10/10 (1.3)	10/10 (1.4)	10/10 (1.3)	
Focal Fibrosis		1/10	6/10	2/10	6/10	3/9	3/10	4/10	6/10	
Inflammation/Necrosis		1/10	1/10	2/10	0/10	0/10	2/10	2/10	2/10	
Diffuse Degeration	–	1/10	1/10	–	–	–	1/10	–		

Shown in parantheses are the mean severity score for cardiomyopathy.