

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Beyond the RfD: Broad Application of a Probabilistic Approach to Improve Chemical Dose–Response Assessments for Noncancer Effects

Weihshueh A. Chiu, Daniel A. Axelrad, Chimeddulam Dalaijamts, Chris Dockins, Kan Shao, Andrew J. Shapiro, and Greg Paoli

Table of Contents

Figure S1. Comparison across sources for Reference Doses (A), Points of Departure (B), and Endpoint-specific Reference Doses (C) for the same chemical evaluated. No comparisons were performed for PPRTVs and HEAST because these sources had very few, if any, (between 0 and 3) overlapping chemicals with other sources. ATSDR = Agency for Toxic Substances and Disease Registry; HEAST = Health Effects Assessment Summary Tables; IRIS = U.S. EPA Integrated Risk Information System; OPP = U.S. EPA Office of Pesticide Programs; PPRTV = Provisional Peer-Reviewed Toxicity Values.

Figure S2. Comparison of “Approximate” Calculation Results with Monte-Carlo Simulation-Based Estimates. The boxplots present the ratio between the “approximate” calculations and Monte Carlo-based results using 10 million samples, for the lower 95% confidence (P05), median (P50), and upper 95% confidence (P95) estimates of the HD_M^I . Each boxplot includes the interquartile range (box), median (line in box), 95% confidence interval (whiskers), and values outside 95% confidence interval (circles).

Figure S3. Comparison of results of traditional and probabilistic dose-response assessment, separated by endpoint. Scatterplot of traditional RfD versus probabilistic RfD, defined as shown in **Figure 1** for $I=1\%$. Solid line denotes equality, and black circles are traditional and probabilistic RfDs, with each panel a subset based on endpoint, as labeled.

Figure S4. Comparison of results of traditional and probabilistic dose-response assessment, separated by Point of Departure (POD). Scatterplot of traditional RfD versus probabilistic RfD, defined as shown in **Figure 1** for $I=1\%$. Solid line denotes equality, and black circles are traditional and probabilistic RfDs, with each panel a subset based on POD, as labeled. BMDL = benchmark dose lower confidence limit; NOAEL = No Observed Adverse Effect Level; LOAEL = Lowest Observed Adverse Effect Level.

Figure S5. Comparison of results of traditional and probabilistic dose-response assessment, separated by decade of assessment. Scatterplot of traditional RfD versus probabilistic RfD, defined as shown in **Figure 1** for $I=1\%$. Solid line denotes equality, and black circles are traditional and probabilistic RfDs, with each panel a subset based on the decade in which the traditional RfD was published, as labeled.

Supplemental Figures




Figure S1. Comparison across sources for Reference Doses (A), Points of Departure (B), and Endpoint-specific Reference Doses (C) for the same chemical evaluated. No comparisons were performed for PPRTVs and HEAST because these sources had very few, if any, (between 0 and 3) overlapping chemicals with other sources. ATSDR = Agency for Toxic

Substances and Disease Registry; HEAST = Health Effects Assessment Summary Tables; IRIS = U.S. EPA Integrated Risk Information System; OPP = U.S. EPA Office of Pesticide Programs; PPRTV = Provisional Peer-Reviewed Toxicity Values.




Figure S2. Comparison of “Approximate” Calculation Results with Monte-Carlo Simulation-Based Estimates. The boxplots present the ratio between the “approximate” calculations and Monte Carlo-based results using 10 million samples, for the lower 95% confidence (P05), median (P50), and upper 95% confidence (P95) estimates of the HD_M^I . Each boxplot includes the interquartile range (box), median (line in box), 95% confidence interval (whiskers), and values outside 95% confidence interval (circles).




Figure S3. Comparison of results of traditional and probabilistic dose-response assessment, separated by endpoint. Scatterplot of traditional RfD versus probabilistic RfD, defined as shown in **Figure 1** for I=1%. Solid line denotes equality, and black circles are traditional and probabilistic RfDs, with each panel a subset based on endpoint, as labeled.




Figure S4. Comparison of results of traditional and probabilistic dose-response assessment, separated by Point of Departure (POD). Scatterplot of traditional RfD versus probabilistic RfD, defined as shown in **Figure 1** for I=1%. Solid line denotes equality, and black circles are traditional and probabilistic RfDs, with each panel a subset based on POD, as labeled. BMDL = benchmark dose lower confidence limit; NOAEL = No Observed Adverse Effect Level; LOAEL = Lowest Observed Adverse Effect Level.




Figure S5. Comparison of results of traditional and probabilistic dose-response assessment, separated by decade of assessment. Scatterplot of traditional RfD versus probabilistic RfD, defined as shown in **Figure 1** for $I=1\%$. Solid line denotes equality, and black circles are traditional and probabilistic RfDs, with each panel a subset based on the decade in which the traditional RfD was published, as labeled.