

### Summary of Estrous Stages and Cycle Data, Bisphenol AF, F2 Rats

Dose ppm	Estrous stage <sup>c</sup> (percent)						Cycle length <sup>a,c,f</sup> (days)			Number of cycles <sup>a,c,f</sup>		
	Proestrus	Estrus	Metestrus	Diestrus	NC <sup>b</sup>	N <sup>d</sup>	Mean	S.E.	N <sup>e</sup>	Mean	S.E.	N <sup>e</sup>
0	5.3	36.6	4.2	53.7	0.2	78	4.88	0.21	71 [18]	2.10	0.10	71 [18]
338	2.1	34.4	3.6	59.3	0.6	77	5.11*	0.19	71 [19]	1.92	0.06	71 [19]
1125	3.1	35.3	1.3	59.7	0.6	20	5.22*	0.17	20 [7]	2.01	0.18	20 [7]

a: Each dose group is compared to the control using the Datta-Satten modified Wilcoxon test with the Hommel adjustment for multiple comparisons. [\*\*=P<.01, \*=P<.05]

b: NC = Not clear, poor quality, or insufficient number of cells

c: Estrous stage percent values are based on overall time in each stage. Mean and S.E. values for cycle length and number of cycles are calculated from the litter means.

d: Number of animals in dose group

e: Number of animals cycling [number of litters]

f: Animals not cycling were excluded from the Cycle Length and Number of Cycles calculations.