Study Number: MOG10866
Test Type: MOG - Range Finding

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

Study Number:

Study Gender:

PWG Approval Date:

Version:

108: Mean Test Compound ConsumptionTest Compound: Isopropylated Phenyl Phosphate

CAS Number: 68937-41-7

MOG10866

Both

See web page for date of PWG Approval

v1.1.7

Date Report Requested: 02/22/2021 Time Report Requested: 07:23:49

Lab: Battelle

Study Number: MOG10866

Days

Test Type: MOG - Range Finding

Route: Dosing in Feed

Phase

Species/Strain: Rat/Harlan Sprague Dawley

0

108: Mean Test Compound Consumption

Test Compound: Isopropylated Phenyl Phosphate

CAS Number: 68937-41-7

Date Report Requested: 02/22/2021 Time Report Requested: 07:23:49

Lab: Battelle

F0 Females

Treatment Groups (ppm) 3000 10000 15000 Dose/Day Dose/Day Dose/Dav (mg/kg/animal/da Ν (mg/kg/animal/da Ν N y) y) y) 12 1328.6 ± 86.8 16 2014.4 ± 172.6 12 12 1656.8 ± 127.3 16 2040.7 ± 233.8 12 16

1000

Study Number: MOG10866

Test Type: MOG - Range Finding

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I08: Mean Test Compound Consumption

Test Compound: Isopropylated Phenyl Phosphate

CAS Number: 68937-41-7

Date Report Requested: 02/22/2021 Time Report Requested: 07:23:49

Lab: Battelle

F0 Females

		Treatment Groups (ppm) 30000	
Phase	Days		
		Dose/Day (mg/kg/animal/da y)	N
Gestation	6 - 9	4150.7 ± 456.5	10
	9 - 12	3246.5 ± 631.0	10
	12 - 15	NR	
	15 - 18	NR	
	18 - 21	NR	
	6 - 21	NR	
Lactation	1 - 4	NR	
	4 - 7	NR	
	7 - 10	NR	
	10 - 14	NR	
	1 - 14	NR	

Study Number: MOG10866
Test Type: MOG - Range Finding

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

108: Mean Test Compound ConsumptionTest Compound: Isopropylated Phenyl Phosphate

CAS Number: 68937-41-7

Date Report Requested: 02/22/2021 Time Report Requested: 07:23:49

Lab: Battelle

LEGEND

Reported as the mean ± SEM. N is the number of animals, number of cages for group housed adult animals or number of litters.

Chemical intake values were excluded when excessive spillage was recorded.

Statistical analysis was not performed on this data.

Consumption is not reported for the non-pregnant animals during gestation and lactation phases

The 30,000 ppm group was terminated on GD12, and the 15,000 ppm group was terminated by LD3/PND3 due to excessive toxicity.

Three dams were removed from the control, 1,000, and 10,000 ppm groups for biological sample collection on GD18, and one pup per sex was selected from each of three dams from the 0 and 1,000 ppm groups on PND4.

NR not recorded

** END OF REPORT **