

Study Number: C11042B-01
Test Type: TOX with Perinatal Exposure
Route: Dosing in Feed
Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption
Test Compound: Triphenyl Phosphate (TPHP)
CAS Number: 115-86-6

Date Report Requested: 05/28/2026
Time Report Requested: 13:56:45
Lab: Battelle

Study Number: C11042B-01
Study Sex: Both
PWG Approval Date: See web page for date of PWG Approval
Version: v1.7.8
Stat Version: 2026.04.15S

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F0 Females

Treatment Groups (ppm)

Phase	Days	0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Gestation	6 - 9	17.4 ± 0.3 **	72.8 ± 1.0 **	23	17.2 ± 0.2	73.0 ± 0.8	31	17.6 ± 0.2	74.8 ± 0.8	30
	9 - 12	18.8 ± 0.3 **	74.5 ± 0.8 **	23	18.5 ± 0.3	74.1 ± 0.9	31	18.6 ± 0.3	74.4 ± 0.9	30
	12 - 15	18.8 ± 0.3 *	69.6 ± 1.0	23	18.3 ± 0.2	68.9 ± 0.7	31	18.5 ± 0.3	69.7 ± 1.0	30
	15 - 18	21.3 ± 0.3 *	71.6 ± 0.8	23	20.8 ± 0.3	70.5 ± 0.8	31	21.1 ± 0.2	72.1 ± 0.7	30
	18 - 21	21.1 ± 0.4	62.7 ± 1.0	20	21.7 ± 0.4	64.3 ± 1.0	28	21.4 ± 0.3	64.7 ± 0.7	27
	6 - 21	19.5 ± 0.3 **	69.4 ± 0.8 *	20	19.4 ± 0.2	69.0 ± 0.7	28	19.4 ± 0.2	70.2 ± 0.7	27
Lactation	1 - 4	36.1 ± 1.3	133.2 ± 3.7	17	32.5 ± 1.0	120.7 ± 3.6	28	33.4 ± 0.9	126.0 ± 3.3	24
	4 - 7	37.6 ± 0.7	135.7 ± 2.3	19	37.8 ± 0.5	135.6 ± 1.9	24	37.3 ± 0.6	137.5 ± 2.2	23
	7 - 10	46.8 ± 0.7 **	164.6 ± 2.0	19	46.3 ± 0.8	162.5 ± 3.1	24	47.2 ± 0.9	169.2 ± 3.0	23
	10 - 13	52.6 ± 0.8 *	179.5 ± 2.2	19	51.0 ± 0.7	173.8 ± 2.6	24	51.3 ± 0.9	178.5 ± 3.0	23
	13 - 16	58.7 ± 1.1 *	198.4 ± 2.8	19	57.7 ± 0.9	191.6 ± 3.1	24	57.4 ± 1.0	195.8 ± 3.1	23
	16 - 19	68.8 ± 2.0	233.5 ± 6.4	19	68.6 ± 1.3	230.2 ± 4.8	24	68.5 ± 1.5	235.5 ± 5.1	23
	19 - 21	87.8 ± 1.8	303.0 ± 6.4	19	87.8 ± 1.8	302.8 ± 6.4	24	88.8 ± 2.0	311.7 ± 7.2	23
	21 - 25	97.6 ± 1.7	346.8 ± 6.3	19	98.2 ± 1.8	349.7 ± 6.4	24	97.9 ± 1.7	355.0 ± 6.7	23
	25 - 28	98.4 ± 1.5	355.7 ± 6.0	19	96.9 ± 1.8	353.5 ± 7.0	24	96.1 ± 1.4	359.2 ± 6.4	23
	1 - 28	65.6 ± 1.0	231.7 ± 2.9	19	64.5 ± 1.0	227.3 ± 3.7	24	64.9 ± 1.0	233.7 ± 3.8	23

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F0 Females

Treatment Groups (ppm)

Phase	Days	300			1000			3000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Gestation	6 - 9	17.8 ± 0.3	74.3 ± 0.7	26	16.5 ± 0.2 *	69.4 ± 0.7 **	29	15.3 ± 0.6 **	65.3 ± 2.3 **	30
	9 - 12	19.0 ± 0.2	74.7 ± 0.7	26	17.8 ± 0.2 *	70.8 ± 0.9 **	29	17.4 ± 0.4 **	70.9 ± 1.6 **	31
	12 - 15	18.6 ± 0.3	68.7 ± 0.8	26	18.2 ± 0.2	68.1 ± 0.7	29	17.9 ± 0.4	68.8 ± 1.4	31
	15 - 18	21.3 ± 0.6	71.4 ± 1.7	26	20.8 ± 0.3	70.9 ± 0.9	28	20.0 ± 0.4	70.3 ± 0.9	31
	18 - 21	21.3 ± 0.5	63.6 ± 1.0	23	20.5 ± 0.3	62.8 ± 0.8	26	20.6 ± 0.4	64.6 ± 0.9	28
	6 - 21	19.5 ± 0.3	69.7 ± 0.6	23	18.8 ± 0.2	67.9 ± 0.5	26	18.2 ± 0.4 *	67.4 ± 0.9	28
Lactation	1 - 4	33.5 ± 1.2	126.0 ± 4.4	22	33.7 ± 1.3	126.0 ± 5.1	25	34.6 ± 1.1	133.8 ± 4.2	27
	4 - 7	37.2 ± 0.4	135.8 ± 2.2	21	35.1 ± 0.8	128.2 ± 3.0	25	36.3 ± 1.0	136.1 ± 3.4	27
	7 - 10	46.8 ± 0.7	167.9 ± 3.7	21	44.4 ± 2.1 *	157.9 ± 7.0	25	43.5 ± 1.2 *	157.9 ± 3.7	27
	10 - 13	52.4 ± 0.7	180.7 ± 2.8	21	47.5 ± 1.5	165.4 ± 5.3	24	47.7 ± 1.5	169.6 ± 4.8	27
	13 - 16	58.9 ± 0.5	198.1 ± 2.5	21	53.4 ± 2.0	182.7 ± 6.8	25	53.9 ± 1.5	188.8 ± 4.7	27
	16 - 19	68.6 ± 1.0	232.5 ± 4.3	21	64.2 ± 2.4	220.9 ± 8.6	25	64.4 ± 1.9	223.3 ± 6.2	27
	19 - 21	88.9 ± 0.9	308.7 ± 5.1	21	83.4 ± 3.5	290.9 ± 12.9	25	81.9 ± 2.9	286.4 ± 9.3	27
	21 - 25	99.5 ± 1.1	355.7 ± 5.6	21	92.8 ± 4.0	330.0 ± 14.8	25	93.4 ± 3.3	335.4 ± 10.9	27
	25 - 28	101.3 ± 1.3	369.4 ± 6.9	21	91.9 ± 3.9	332.7 ± 14.6	25	94.9 ± 3.5	346.4 ± 11.8	27
	1 - 28	65.7 ± 0.5	233.9 ± 3.3	21	61.2 ± 2.2	218.0 ± 8.1	25	61.6 ± 1.9	223.3 ± 6.2	27

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Males: Biosample Cohort

Treatment Groups (ppm)

Phase	Days	0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	16.1 ± 0.4	140.7 ± 3.7	3	13.7 ± 0.2 *	125.7 ± 2.7	3	15.0 ± 0.4	137.0 ± 3.9	3
	35 - 41	20.1 ± 1.3	146.0 ± 2.7	3	18.1 ± 0.3	139.7 ± 2.1	3	18.9 ± 0.7	145.5 ± 1.3	3
	41 - 49	24.9 ± 1.4	120.1 ± 2.3 *	3	23.1 ± 0.3	117.4 ± 0.5	3	23.9 ± 0.8	122.1 ± 4.8	3
	49 - 56	26.3 ± 1.3	103.9 ± 1.9 *	3	25.2 ± 0.1	103.7 ± 1.4	3	26.1 ± 1.4	108.9 ± 6.1	3
	56 - 63	27.3 ± 1.1	93.8 ± 1.1	3	25.5 ± 1.5	90.6 ± 4.1	3	28.3 ± 2.3	102.1 ± 6.8	3
	63 - 70	28.2 ± 1.7	87.5 ± 3.2	3	25.3 ± 1.8	81.9 ± 4.2	3	28.6 ± 2.8	93.7 ± 8.1	3
	70 - 77	29.0 ± 2.0	83.7 ± 4.0	3	26.0 ± 2.2	78.1 ± 4.4	3	28.8 ± 0.9	88.2 ± 1.4	3
	77 - 84	29.1 ± 2.1	79.6 ± 3.8	3	25.5 ± 2.0	71.9 ± 3.5	3	27.8 ± 2.4	80.9 ± 5.8	3
	84 - 91	29.5 ± 2.6	77.4 ± 5.0	3	24.7 ± 2.1	66.3 ± 3.4	3	27.0 ± 2.4	75.5 ± 5.9	3
	91 - 98	26.5 ± 0.1	69.1 ± 0.8	2	24.1 ± 2.1	63.0 ± 2.9	3	26.3 ± 2.0	71.7 ± 5.3	3
	98 - 105	27.8 ± 2.3	69.1 ± 4.1	3	23.9 ± 1.7	60.7 ± 2.2	3	24.9 ± 1.7	66.2 ± 4.0	3
	105 - 112	25.4 ± 0.2	63.2 ± 1.5 *	2	24.1 ± 2.3	59.5 ± 3.3	3	24.9 ± 1.7	64.7 ± 4.2	3
	112 - 119	23.7 ± 1.1	57.9 ± 1.9	2	22.9 ± 2.5	55.5 ± 3.6	3	24.3 ± 1.7	62.0 ± 4.0	3
	119 - 126	25.4 ± 0.8	60.6 ± 0.7	2	24.5 ± 2.6	58.4 ± 4.0	3	25.1 ± 3.4	64.5 ± 8.1	2
28 - 126	26.0 ± 1.4	81.9 ± 2.2	3	23.5 ± 1.5	76.2 ± 2.6	3	24.6 ± 2.6	84.2 ± 7.9	2	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Males: Biosample Cohort

Treatment Groups (ppm)

Phase	Days	300			1000			3000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	15.4 ± 0.4	141.7 ± 4.8	3	15.0 ± 0.4	138.7 ± 3.5	3	13.9 ± 0.5	135.8 ± 7.5	3
	35 - 41	19.1 ± 0.6	147.7 ± 1.7	3	19.0 ± 0.9	150.0 ± 8.4	3	19.9 ± 1.4	166.9 ± 22.1	3
	41 - 49	25.6 ± 1.9	131.4 ± 10.3	3	23.2 ± 0.5	122.1 ± 1.0	3	23.6 ± 0.4	132.1 ± 8.9	3
	49 - 56	28.3 ± 2.6	119.1 ± 11.6	3	25.0 ± 0.7	106.9 ± 1.3	3	26.0 ± 1.0	118.0 ± 5.3	3
	56 - 63	27.4 ± 1.6	100.2 ± 6.7	3	25.1 ± 0.1	93.3 ± 1.3	3	25.4 ± 1.1	101.6 ± 9.2	2
	63 - 70	27.7 ± 2.8	91.9 ± 10.0	3	24.2 ± 0.3	82.3 ± 2.1	3	25.4 ± 0.9	91.8 ± 7.2	2
	70 - 77	27.6 ± 2.6	85.5 ± 9.0	3	25.2 ± 0.2	79.6 ± 0.7	3	26.7 ± 1.8	88.4 ± 5.4	3
	77 - 84	26.8 ± 2.5	78.6 ± 7.9	3	26.3 ± 0.5	77.3 ± 0.8	3	25.2 ± 0.5	79.3 ± 1.6	2
	84 - 91	23.4 ± 0.6	65.3 ± 0.1	2	25.5 ± 0.3	71.1 ± 0.3	3	24.6 ± 0.6	72.9 ± 0.2	3
	91 - 98	27.5 ± 3.2	74.9 ± 8.8	3	25.8 ± 0.3	69.6 ± 1.0	3	24.3 ± 0.5	69.7 ± 1.8	3
	98 - 105	24.2 ± 0.4	64.1 ± 0.2	2	26.1 ± 0.7	68.2 ± 0.7	3	25.0 ± 0.7	69.4 ± 2.1	3
	105 - 112	30.0 ± 5.5	77.4 ± 14.4	3	25.1 ± 0.4	64.1 ± 1.1	3	25.5 ± 0.9	69.2 ± 1.2	3
	112 - 119	24.6 ± 0.6	62.3 ± 1.2	3	23.5 ± 0.3	58.9 ± 0.5	3	23.6 ± 0.6	62.7 ± 1.4	3
	119 - 126	30.6 ± 5.6	74.8 ± 13.2	3	24.8 ± 0.4	61.1 ± 1.4	3	23.8 ± 0.5	61.7 ± 1.0	3
28 - 126	25.7 ± 2.0	86.0 ± 7.1	3	23.9 ± 0.3	80.4 ± 0.4	3	23.6 ± 0.1	84.0 ± 2.6	3	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Females: Biosample Cohort

Treatment Groups (ppm)

Phase	Days	0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	12.9 ± 0.8	134.3 ± 2.4 **	2	13.3 ± 0.5	137.3 ± 1.4	2	13.5 ± 0.8	138.5 ± 0.9	2
	35 - 41	15.2 ± 1.0	135.9 ± 4.1 *	2	15.6 ± 0.2	139.1 ± 1.6	2	17.3 ± 0.9	152.2 ± 1.0	2
	41 - 49	18.9 ± 1.3	123.0 ± 4.3	2	19.8 ± 1.3	129.0 ± 5.3	2	19.0 ± 0.6	124.0 ± 0.1	2
	49 - 56	21.6 ± 1.9	123.7 ± 13.5	2	20.3 ± 0.0	117.7 ± 2.6	2	20.2 ± 2.4	116.8 ± 10.1	2
	56 - 63	20.5 ± 1.5	105.1 ± 7.1	2	21.5 ± 2.9	113.5 ± 12.1	2	19.2 ± 1.7	101.4 ± 5.6	2
	63 - 70	21.5 ± 1.0	102.7 ± 4.2	2	20.5 ± 2.9	103.3 ± 12.2	2	19.3 ± 2.2	95.0 ± 8.3	2
	70 - 77	21.7 ± 1.6 *	98.6 ± 7.5	2	20.9 ± 2.0	100.8 ± 8.0	2	19.5 ± 2.1	91.1 ± 7.4	2
	77 - 84	23.0 ± 1.8	99.9 ± 8.2	2	19.5 ± 0.3	90.0 ± 2.7	2	21.0 ± 4.6	93.8 ± 18.3	2
	84 - 91	20.4 ± 1.1	85.0 ± 4.1	2	19.3 ± 0.3	85.9 ± 0.0	2	18.0 ± 1.6	78.6 ± 4.8	2
	91 - 98	22.9	95.0	1	18.9 ± 1.5	82.3 ± 5.7	2	18.0 ± 1.6	77.2 ± 5.0	2
	98 - 105	17.8 ± 1.1	72.4 ± 4.7	2	18.6 ± 0.7	80.1 ± 2.2	2	17.4 ± 1.6	73.8 ± 5.0	2
	105 - 112	17.6 ± 0.3	70.0 ± 0.7	2	18.5 ± 0.7	78.7 ± 3.0	2	16.8 ± 0.9	69.7 ± 1.8	2
	112 - 119	16.6 ± 0.1	66.0 ± 0.1	2	16.6 ± 1.2	70.7 ± 5.3	2	15.8 ± 1.3	65.5 ± 3.7	2
	119 - 126	17.7 ± 0.5	70.0 ± 2.3	2	17.8 ± 1.1	75.2 ± 4.3	2	17.0 ± 1.9	69.6 ± 5.7	2
28 - 126	19.0 ± 0.2	92.9 ± 0.3	2	18.7 ± 1.1	96.4 ± 4.1	2	18.1 ± 1.8	91.2 ± 6.2	2	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Females: Biosample Cohort

Treatment Groups (ppm)

Phase	Days	300			1000			3000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	12.5 ± 0.3	139.2 ± 0.4	2	13.7 ± 0.6	140.5 ± 7.3	3	13.3 ± 0.7	144.3 ± 1.5 *	3
	35 - 41	15.3 ± 0.4	147.9 ± 7.1	2	16.4 ± 0.1	144.3 ± 2.0	3	19.1 ± 2.8	175.8 ± 27.8	3
	41 - 49	18.7 ± 2.0	128.4 ± 12.6	2	19.1 ± 1.0	122.7 ± 4.8	2	16.5 ± 0.9	118.9 ± 9.1	2
	49 - 56	18.8 ± 1.1	111.8 ± 5.7	2	24.1 ± 2.7	138.8 ± 15.9	3	17.3 ± 0.7	110.0 ± 7.4	2
	56 - 63	18.9 ± 1.8	101.1 ± 9.0	2	21.7 ± 1.5	114.0 ± 7.4	3	17.9 ± 1.2	103.5 ± 9.2	2
	63 - 70	18.6 ± 1.7	93.3 ± 7.9	2	25.0 ± 1.7	122.5 ± 7.6	3	20.4 ± 3.4	106.2 ± 14.4	3
	70 - 77	18.2 ± 2.4	87.2 ± 10.9	2	20.1 ± 1.4	94.4 ± 6.6	3	15.9 ± 0.3	82.5 ± 4.9	2
	77 - 84	18.4 ± 2.1	84.4 ± 8.6	2	21.6 ± 1.7	98.5 ± 6.0	3	16.1 ± 0.7	80.2 ± 0.8	2
	84 - 91	18.0 ± 0.7	80.0 ± 3.0	2	24.1 ± 2.8	108.1 ± 15.6	2	17.6 ± 0.5	82.7 ± 2.2	3
	91 - 98	17.4 ± 2.0	75.7 ± 8.7	2	17.8 ± 1.1	76.0 ± 3.8	3	15.7 ± 0.8	72.4 ± 5.9	3
	98 - 105	17.7 ± 1.6	75.2 ± 6.7	2	17.2 ± 0.3	72.8 ± 0.7	2	19.1 ± 2.0	85.8 ± 6.9	3
	105 - 112	17.7 ± 0.9	73.3 ± 3.9	2	18.3 ± 1.1	75.6 ± 3.9	3	16.2 ± 0.1	74.4 ± 4.1	2
	112 - 119	17.1 ± 2.2	70.0 ± 8.9	2	18.7 ± 3.1	76.6 ± 11.9	3	15.0 ± 0.7	66.2 ± 1.6	3
	119 - 126	19.2 ± 3.7	77.8 ± 14.3	2	18.4 ± 1.5	74.5 ± 5.5	3	16.0 ± 0.3	71.4 ± 3.1	3
28 - 126	17.7 ± 1.6	90.7 ± 7.8	2	20.0 ± 1.1	100.8 ± 5.5	3	16.7 ± 0.4	90.5 ± 2.1	3	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Males: Subchronic Cohort

Treatment Groups (ppm)

Phase	Days	0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	15.2 ± 0.2	135.9 ± 1.7	6	15.1 ± 0.2	136.0 ± 3.8	6	15.3 ± 0.6	140.0 ± 3.4	6
	35 - 41	20.4 ± 0.9	153.0 ± 5.6	6	20.2 ± 0.5	154.3 ± 5.3	6	19.7 ± 0.9	152.3 ± 3.6	6
	41 - 49	24.8 ± 0.5	124.0 ± 3.3 *	6	23.7 ± 0.6	118.9 ± 2.7	6	23.7 ± 0.8	121.9 ± 2.0	6
	49 - 56	25.2 ± 1.2	105.0 ± 3.3 **	6	25.4 ± 0.7	104.1 ± 2.3	6	25.6 ± 1.2	107.4 ± 3.3	6
	56 - 63	27.0 ± 0.9	99.2 ± 6.2 *	6	26.2 ± 0.6	93.3 ± 1.3	6	26.0 ± 1.3	94.3 ± 3.3	6
	63 - 70	26.7 ± 0.7	87.6 ± 2.4	6	26.4 ± 0.7	85.6 ± 1.5	6	26.6 ± 1.6	87.9 ± 4.1	6
	70 - 77	26.8 ± 0.6	81.5 ± 1.9	6	25.8 ± 0.6	78.2 ± 1.5	6	26.3 ± 1.6	81.2 ± 3.8	6
	77 - 84	26.7 ± 0.6	76.8 ± 1.7	6	26.5 ± 0.7	76.2 ± 1.3	6	26.3 ± 1.4	77.2 ± 3.0	6
	84 - 91	26.1 ± 0.6	72.4 ± 1.6	6	25.8 ± 0.6	71.1 ± 1.9	6	25.7 ± 1.2	72.4 ± 2.5	6
	91 - 98	25.7 ± 0.6	69.2 ± 1.3	6	25.7 ± 0.7	68.6 ± 2.0	6	26.5 ± 1.6	71.9 ± 2.6	6
	98 - 105	24.5 ± 0.4	64.2 ± 0.9	6	25.6 ± 0.7	66.6 ± 1.9	6	23.7 ± 0.8	64.4 ± 1.8	5
	105 - 112	24.2 ± 0.7	61.9 ± 1.5 *	6	25.2 ± 1.0	63.8 ± 2.7	6	24.1 ± 1.2	62.1 ± 1.7	6
	112 - 119	24.1 ± 0.5	60.5 ± 1.0	6	24.7 ± 0.9	61.1 ± 2.5	6	24.0 ± 1.2	60.5 ± 1.5	6
28 - 119	24.4 ± 0.3	83.2 ± 1.1	6	24.4 ± 0.5	82.3 ± 1.6	6	24.2 ± 1.1	83.4 ± 2.2	6	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Males: Subchronic Cohort

Treatment Groups (ppm)

Phase	Days	300			1000			3000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	15.2 ± 0.4	142.9 ± 3.8	6	14.9 ± 0.4	136.4 ± 2.5	6	14.5 ± 0.3	142.4 ± 2.3	6
	35 - 41	19.8 ± 0.6	155.5 ± 5.9	6	20.0 ± 0.8	156.5 ± 5.0	6	19.5 ± 1.2	162.6 ± 7.9	6
	41 - 49	25.1 ± 1.2	128.3 ± 6.3	6	25.6 ± 1.4	132.9 ± 6.1	6	25.1 ± 1.8	132.8 ± 5.0	6
	49 - 56	26.1 ± 0.9	109.4 ± 3.9	6	27.4 ± 1.4	116.0 ± 4.3	6	26.0 ± 1.1	116.1 ± 2.8	6
	56 - 63	27.5 ± 1.0	99.6 ± 3.5	6	27.4 ± 1.1	100.1 ± 2.7	6	26.4 ± 1.2	102.9 ± 2.8	6
	63 - 70	28.4 ± 1.5	92.3 ± 4.4	6	27.9 ± 1.8	91.8 ± 4.1	6	25.6 ± 1.1	90.4 ± 1.9	6
	70 - 77	28.7 ± 1.0	87.1 ± 2.5	6	27.1 ± 1.6	83.1 ± 3.7	6	24.6 ± 1.1	81.4 ± 1.6	6
	77 - 84	27.5 ± 1.1	79.3 ± 2.8	6	26.8 ± 1.2	78.2 ± 2.1	6	25.2 ± 1.6	79.4 ± 3.3	6
	84 - 91	27.2 ± 1.1	74.9 ± 2.6	6	25.9 ± 1.1	72.3 ± 1.7	6	23.6 ± 0.9	71.7 ± 0.8	6
	91 - 98	25.9 ± 0.7	70.1 ± 2.1	5	25.8 ± 0.6	69.8 ± 1.3	6	23.8 ± 0.9	69.9 ± 1.3	6
	98 - 105	25.2 ± 1.1	66.6 ± 3.1	5	25.6 ± 0.9	67.2 ± 2.1	6	23.3 ± 0.6	66.8 ± 2.0	6
	105 - 112	26.6 ± 0.6	67.3 ± 1.1	6	24.9 ± 1.0	63.7 ± 1.5	6	23.3 ± 0.9	64.7 ± 1.0	6
	112 - 119	26.7 ± 0.8	66.1 ± 1.7	6	24.2 ± 0.8	60.6 ± 1.7	6	22.7 ± 0.9	61.8 ± 1.2	6
28 - 119	25.6 ± 0.8	86.9 ± 2.4	6	24.9 ± 0.9	85.4 ± 1.8	6	23.4 ± 0.8	86.5 ± 1.3	6	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Females: Subchronic Cohort

Treatment Groups (ppm)

Phase	Days	0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	12.4 ± 0.3	133.8 ± 3.3	4	12.4 ± 0.2	127.0 ± 1.0	3	12.5 ± 0.3	129.7 ± 2.5	4
	35 - 41	17.3 ± 2.4	159.6 ± 20.2	4	15.3 ± 0.2	136.4 ± 3.2	3	15.3 ± 0.6	138.1 ± 6.3	4
	41 - 49	18.5 ± 0.7	123.2 ± 2.9	4	17.7 ± 0.4	115.0 ± 5.2	3	17.7 ± 0.6	116.0 ± 2.8	4
	49 - 56	18.4 ± 0.8	107.5 ± 4.4	4	18.5 ± 1.2	105.8 ± 9.1	3	18.7 ± 0.3	106.7 ± 0.7	4
	56 - 63	18.2 ± 0.8	96.7 ± 3.8	4	18.1 ± 0.8	94.1 ± 5.8	3	18.8 ± 0.7	97.8 ± 3.8	4
	63 - 70	17.9 ± 0.4	89.1 ± 1.9 *	4	17.0 ± 0.7	82.8 ± 3.0	3	19.2 ± 1.2	93.4 ± 5.0	4
	70 - 77	17.3 ± 0.5	81.6 ± 1.5	4	16.7 ± 0.2	77.9 ± 2.9	3	19.6 ± 0.5	90.2 ± 1.1	4
	77 - 84	19.8 ± 1.3	89.6 ± 6.4	4	16.5 ± 0.8	73.9 ± 2.7	3	19.5 ± 0.8	86.7 ± 2.1	4
	84 - 91	17.7 ± 0.5 *	77.3 ± 0.3	4	16.5 ± 0.4	72.3 ± 2.0	3	18.4 ± 0.8	79.3 ± 4.0	4
	91 - 98	15.9 ± 0.5	68.9 ± 1.5	4	17.4 ± 0.6	74.0 ± 5.6	3	17.2 ± 0.6	72.7 ± 2.6	4
	98 - 105	16.6 ± 1.0	71.2 ± 4.0	4	15.5 ± 0.1	64.2 ± 1.5	3	18.0 ± 1.0	74.6 ± 4.6	4
	105 - 112	15.7 ± 0.5	65.9 ± 1.1	4	15.5 ± 0.3	63.7 ± 1.9	3	16.9 ± 0.6	68.9 ± 2.1	4
	112 - 119	14.5 ± 0.3	61.1 ± 1.3	4	15.2 ± 0.5	61.5 ± 1.1	3	14.2 ± 1.7	57.5 ± 6.3	4
28 - 119	17.0 ± 0.5	88.7 ± 2.4	4	16.4 ± 0.1	83.4 ± 2.6	3	17.4 ± 0.4	88.5 ± 1.1	4	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Females: Subchronic Cohort

Treatment Groups (ppm)

Phase	Days	300			1000			3000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	12.9 ± 0.5	135.6 ± 2.9	4	12.7 ± 0.3	133.3 ± 3.6	3	12.2 ± 0.6	135.6 ± 6.3	4
	35 - 41	15.5 ± 1.0	141.8 ± 5.6	4	15.9 ± 1.0	145.6 ± 10.7	3	16.0 ± 1.4	156.2 ± 13.9	4
	41 - 49	18.2 ± 1.3	121.3 ± 7.7	4	17.8 ± 1.0	119.0 ± 9.0	3	17.6 ± 0.8	124.7 ± 6.2	4
	49 - 56	19.0 ± 1.1	110.2 ± 5.2	4	18.3 ± 0.9	106.9 ± 6.7	3	18.7 ± 0.9	114.8 ± 5.8	4
	56 - 63	18.3 ± 1.0	97.1 ± 5.0	4	17.5 ± 0.7	93.1 ± 4.9	3	18.6 ± 1.0	103.6 ± 6.2	4
	63 - 70	19.7 ± 0.7	98.1 ± 3.1	4	18.2 ± 0.4	90.5 ± 2.8	3	18.9 ± 0.9	97.6 ± 5.7	4
	70 - 77	17.4 ± 0.7	82.4 ± 3.0	4	17.7 ± 1.0	83.7 ± 6.2	3	17.3 ± 0.3	84.6 ± 1.6	4
	77 - 84	17.2 ± 0.8	79.0 ± 4.0	4	17.7 ± 0.6	80.6 ± 4.2	3	18.6 ± 0.9	87.3 ± 4.4	4
	84 - 91	16.5 ± 0.3	73.4 ± 1.6	4	17.1 ± 1.3	75.4 ± 6.6	3	16.0 ± 0.3	73.1 ± 0.8	4
	91 - 98	16.3 ± 0.5	71.2 ± 2.0	4	15.5 ± 1.9	67.6 ± 8.7	3	16.6 ± 0.6	74.0 ± 3.8	4
	98 - 105	16.2 ± 0.7	69.4 ± 3.3	4	17.9 ± 0.9	76.9 ± 4.0	3	15.6 ± 0.7	68.8 ± 2.7	4
	105 - 112	15.4 ± 0.4	65.4 ± 2.1	4	16.8 ± 0.8	70.1 ± 3.4	3	15.9 ± 0.8	69.6 ± 4.1	4
	112 - 119	14.7 ± 0.2	61.7 ± 1.7	4	15.5 ± 0.5	64.0 ± 2.9	3	14.4 ± 0.3	62.2 ± 1.2	4
28 - 119	16.8 ± 0.5	87.8 ± 2.5	4	16.8 ± 0.7	87.7 ± 4.7	3	16.7 ± 0.4	90.4 ± 2.9	4	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Males: Neurobehavioral Cohort

Treatment Groups (ppm)

Phase	Days	0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	13.7 ± 0.3	121.2 ± 1.3 *	10	14.0 ± 0.2	123.7 ± 2.3	11	13.8 ± 0.1	124.0 ± 1.6	11
	35 - 41	17.2 ± 0.5	126.1 ± 1.9 **	10	17.8 ± 0.2	130.1 ± 1.9	11	17.4 ± 0.3	129.2 ± 1.6	11
	41 - 49	21.8 ± 0.6	102.6 ± 1.3 **	10	22.2 ± 0.3	103.5 ± 1.2	11	22.1 ± 0.3	104.7 ± 0.9	11
	49 - 56	23.8 ± 0.7	91.3 ± 1.3 **	10	23.9 ± 0.3	92.4 ± 1.0	11	24.3 ± 0.3	94.7 ± 0.5 *	11
	56 - 63	23.8 ± 0.7	80.0 ± 1.2 *	10	24.0 ± 0.4	82.0 ± 1.4	11	24.4 ± 0.4	83.5 ± 1.0	11
	63 - 70	24.3 ± 0.7	74.8 ± 1.3 *	10	23.9 ± 0.5	75.4 ± 1.1	11	24.7 ± 0.3	77.0 ± 0.7	11
	70 - 77	22.6 ± 0.6	66.0 ± 1.2	10	22.9 ± 1.0	68.4 ± 2.5	11	24.2 ± 0.5	70.7 ± 1.1	11
	77 - 84	24.2 ± 0.9	67.1 ± 3.0	10	23.2 ± 0.6	66.2 ± 1.1	11	24.2 ± 0.5	67.4 ± 0.9	11
	84 - 91	25.3 ± 0.5	66.8 ± 0.8	10	24.9 ± 0.5	67.7 ± 0.9	11	24.0 ± 0.7	64.5 ± 1.3	11
	91 - 98	25.2 ± 0.6	63.6 ± 0.7	10	24.4 ± 0.4	63.6 ± 0.9	11	23.7 ± 0.7	61.9 ± 1.2	11
	98 - 105	24.6 ± 0.6 *	60.1 ± 0.9	10	23.8 ± 0.5	60.1 ± 1.2	11	23.9 ± 0.7	60.8 ± 1.5	11
	105 - 112	23.9 ± 0.5 **	57.0 ± 1.3	10	23.1 ± 0.6	57.0 ± 1.3	11	23.1 ± 0.6	57.2 ± 1.2	11
	112 - 119	23.2 ± 0.6 *	54.2 ± 0.7	10	22.8 ± 0.5	55.2 ± 1.4	11	23.1 ± 0.6	56.3 ± 1.1	11
	119 - 126	23.5 ± 0.7 *	53.9 ± 0.9	10	22.9 ± 0.6	54.6 ± 1.4	11	23.1 ± 0.7	55.1 ± 1.2	11
126 - 133	22.3 ± 0.7	50.1 ± 1.3	10	22.8 ± 0.6	53.0 ± 1.3	11	22.9 ± 0.7	53.8 ± 1.4	11	
28 - 133	22.6 ± 0.4	68.6 ± 0.9	10	22.5 ± 0.4	70.4 ± 1.0	11	22.7 ± 0.4	70.9 ± 0.9	11	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Males: Neurobehavioral Cohort

Treatment Groups (ppm)

Phase	Days	300			1000			3000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	13.8 ± 0.2	122.5 ± 1.1	11	14.2 ± 0.3	124.6 ± 1.4	11	13.3 ± 0.2	127.2 ± 1.7 *	11
	35 - 41	17.7 ± 0.3	130.3 ± 1.9	11	18.0 ± 0.3	131.5 ± 1.7	11	17.2 ± 0.4	136.3 ± 2.2 **	11
	41 - 49	22.3 ± 0.3	105.8 ± 0.8 *	11	22.7 ± 0.3	106.0 ± 0.9 *	11	22.1 ± 0.4	111.1 ± 1.5 **	11
	49 - 56	24.5 ± 0.3	95.0 ± 1.5 *	11	24.7 ± 0.2	94.4 ± 0.6 *	11	23.7 ± 0.2	97.8 ± 1.5 **	11
	56 - 63	24.8 ± 0.3	83.9 ± 0.9	11	24.5 ± 0.3	82.5 ± 0.8	11	23.7 ± 0.4	85.3 ± 1.7 *	11
	63 - 70	25.2 ± 0.3	77.7 ± 0.9	11	24.8 ± 0.3	76.7 ± 1.0	11	23.8 ± 0.6	78.4 ± 2.0	11
	70 - 77	25.2 ± 0.5 *	72.5 ± 1.5 *	11	22.8 ± 0.4	66.7 ± 0.9	11	22.2 ± 0.5	69.0 ± 1.2	11
	77 - 84	23.2 ± 1.3	64.3 ± 3.2	11	23.7 ± 0.4	66.5 ± 0.7	11	22.9 ± 0.5	68.0 ± 1.5	11
	84 - 91	24.6 ± 0.6	66.3 ± 2.5	11	25.6 ± 0.5	68.5 ± 0.9	11	24.2 ± 0.4	68.6 ± 0.8	11
	91 - 98	24.0 ± 0.3	62.0 ± 0.9	11	24.7 ± 0.5	63.9 ± 0.9	11	23.7 ± 0.4	64.5 ± 0.8	11
	98 - 105	23.8 ± 0.4	59.8 ± 0.8	11	23.7 ± 0.5	60.0 ± 0.8	11	22.7 ± 0.4	60.1 ± 1.0	11
	105 - 112	22.9 ± 0.3	56.3 ± 0.7	11	22.5 ± 0.6	55.8 ± 1.1	11	21.6 ± 0.3 **	55.9 ± 0.8	11
	112 - 119	23.0 ± 0.4	55.5 ± 0.9	11	22.3 ± 0.5	54.5 ± 0.8	11	21.4 ± 0.3	54.6 ± 1.0	11
	119 - 126	22.6 ± 0.4	53.7 ± 0.8	11	22.6 ± 0.4	54.1 ± 0.8	11	21.7 ± 0.4	54.4 ± 0.9	11
126 - 133	22.8 ± 0.5	53.2 ± 0.9	11	22.7 ± 0.5	53.1 ± 1.0	11	21.4 ± 0.4	52.6 ± 0.8	11	
28 - 133	22.8 ± 0.3	70.7 ± 0.7	11	22.8 ± 0.3	70.8 ± 0.6	11	21.8 ± 0.3	71.8 ± 1.0	11	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Females: Neurobehavioral Cohort

Treatment Groups (ppm)

Phase	Days	0			30			100		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	11.6 ± 0.6	115.7 ± 3.3 **	6	11.1 ± 0.3	112.3 ± 2.2	6	11.0 ± 0.2	115.4 ± 2.4	6
	35 - 41	13.6 ± 0.7	116.3 ± 3.4	6	13.5 ± 0.2	116.3 ± 2.0	6	13.3 ± 0.3	118.5 ± 3.3	6
	41 - 49	16.6 ± 0.6	102.3 ± 2.3	6	16.5 ± 0.7	104.1 ± 5.7	6	16.3 ± 0.7	105.3 ± 5.0	6
	49 - 56	17.1 ± 0.7	92.6 ± 2.3	6	17.4 ± 0.5	95.6 ± 3.3	6	16.9 ± 0.4	94.9 ± 2.5	6
	56 - 63	16.0 ± 0.4 *	79.5 ± 0.5	6	16.2 ± 0.4	80.9 ± 2.4	6	16.3 ± 0.3	82.9 ± 1.7	6
	63 - 70	16.9 ± 0.8	78.6 ± 2.7	6	16.2 ± 0.5	76.4 ± 2.4	6	16.3 ± 0.5	78.2 ± 2.5	6
	70 - 77	16.2 ± 0.7	71.3 ± 1.6	6	15.3 ± 0.6	68.4 ± 2.0	6	16.6 ± 0.6	75.8 ± 2.9	6
	77 - 84	16.7 ± 0.5 *	71.3 ± 1.3	6	15.8 ± 0.4	67.9 ± 0.7	6	17.0 ± 0.8	74.3 ± 3.0	6
	84 - 91	17.3 ± 0.3	71.4 ± 0.7	6	17.0 ± 0.5	70.4 ± 2.1	6	16.1 ± 0.5	67.9 ± 1.8	6
	91 - 98	16.7 ± 0.5	67.0 ± 1.4	6	16.6 ± 0.4	66.8 ± 1.6	6	15.6 ± 0.5	64.8 ± 2.2	6
	98 - 105	15.9 ± 0.5	62.8 ± 1.3	6	16.1 ± 0.6	63.6 ± 1.8	6	15.9 ± 0.4	65.2 ± 1.7	6
	105 - 112	15.7 ± 0.7	61.1 ± 1.9	6	15.5 ± 0.5	60.2 ± 1.3	6	15.5 ± 0.4	62.1 ± 1.6	6
	112 - 119	15.4 ± 0.4	59.4 ± 1.2	6	14.6 ± 0.4	56.3 ± 1.3	6	14.6 ± 0.4	58.1 ± 1.7	6
	119 - 126	15.4 ± 0.3	58.4 ± 1.0	6	15.0 ± 0.4	57.4 ± 1.0	6	14.6 ± 0.4	57.4 ± 1.5	6
126 - 133	15.7 ± 0.5	59.0 ± 1.8	6	14.8 ± 0.6	55.9 ± 1.7	6	14.8 ± 0.4	57.6 ± 1.0	6	
28 - 133	15.9 ± 0.4	74.2 ± 1.1	6	15.5 ± 0.4	73.0 ± 1.4	6	15.5 ± 0.3	74.8 ± 1.6	6	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

F1 Females: Neurobehavioral Cohort

Treatment Groups (ppm)

Phase	Days	300			1000			3000		
		Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N	Wt (g/animal/day)	Wt (g/kg/day)	N
Postnatal	28 - 35	11.5 ± 0.2	117.1 ± 1.5	7	11.4 ± 0.2	117.4 ± 2.1	7	11.8 ± 0.4	124.5 ± 2.6 *	7
	35 - 41	13.6 ± 0.3	117.1 ± 1.9	7	13.8 ± 0.4	120.8 ± 4.0	7	13.7 ± 0.4	123.5 ± 2.9	7
	41 - 49	16.5 ± 0.6	103.7 ± 3.0	7	16.6 ± 1.3	107.1 ± 8.3	7	15.7 ± 0.4	102.6 ± 1.6	7
	49 - 56	17.5 ± 0.7	96.6 ± 2.9	7	16.3 ± 0.4	92.9 ± 1.9	7	16.2 ± 0.4	93.2 ± 1.6	7
	56 - 63	16.9 ± 0.7	84.7 ± 2.4	7	15.3 ± 0.4	80.1 ± 1.7	7	15.0 ± 0.3	79.8 ± 1.2	7
	63 - 70	16.8 ± 0.6	79.5 ± 2.0	7	15.7 ± 0.5	77.4 ± 2.0	7	15.6 ± 0.4	78.1 ± 0.9	7
	70 - 77	15.9 ± 0.3	71.8 ± 0.8	7	14.5 ± 0.4	68.3 ± 1.7	7	14.9 ± 0.5	70.7 ± 1.4	7
	77 - 84	16.3 ± 0.5	70.7 ± 1.6	7	15.2 ± 0.3	69.2 ± 1.4	7	15.6 ± 0.5	70.8 ± 1.7	7
	84 - 91	16.1 ± 0.5	68.0 ± 2.0	7	16.6 ± 0.4	72.7 ± 1.7	7	16.6 ± 0.3	73.0 ± 1.1	7
	91 - 98	16.1 ± 0.4	66.6 ± 1.0	7	15.3 ± 0.5	65.5 ± 2.2	7	16.1 ± 0.6	68.6 ± 1.6	7
	98 - 105	15.9 ± 0.5	64.7 ± 1.5	7	15.6 ± 0.5	65.2 ± 1.8	7	15.6 ± 0.5	65.6 ± 1.3	7
	105 - 112	15.4 ± 0.4	61.8 ± 1.1	7	14.9 ± 0.6	61.2 ± 2.2	7	14.8 ± 0.7	61.4 ± 2.2	7
	112 - 119	15.3 ± 0.5	60.7 ± 1.8	7	14.6 ± 0.5	59.5 ± 1.5	7	14.9 ± 0.6	61.1 ± 1.7	7
	119 - 126	15.2 ± 0.4	59.6 ± 1.0	7	14.8 ± 0.7	59.1 ± 2.2	7	14.6 ± 0.5	59.5 ± 1.3	7
126 - 133	14.9 ± 0.4	57.0 ± 1.3	7	14.8 ± 0.6	58.2 ± 1.7	7	14.8 ± 0.6	59.1 ± 1.5	7	
28 - 133	15.6 ± 0.4	74.8 ± 1.3	7	15.1 ± 0.4	74.5 ± 1.9	7	15.1 ± 0.4	75.4 ± 1.1	7	

Study Number: C11042B-01

Test Type: TOX with Perinatal Exposure

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

I06: Mean Feed Consumption

Test Compound: Triphenyl Phosphate (TPHP)

CAS Number: 115-86-6

Date Report Requested: 05/28/2026

Time Report Requested: 13:56:45

Lab: Battelle

LEGEND

Reported as the mean \pm SEM. N is the number of animals, or number of cages for group housed adult animals.

F0 females were singly housed. All F1 cohorts were group housed.

Feed consumption values were excluded when excessive spillage was recorded.

Consumption is not reported for the non-pregnant animals during gestation and lactation phases.

Consumption data for group housed adult animals in the F1 generation is analyzed with cage as the statistical unit.

Group housed animals were housed up to four animals per cage.

Statistical analysis was performed by Jonckheere (trend) and Shirley or Dunn (pairwise) tests.

Statistical significance for the control group indicates a significant trend test.

Statistical significance for a treatment group indicates a significant pairwise test compared to the vehicle control group.

* Statistically significant at $p \leq 0.05$

** Statistically significant at $p \leq 0.01$

SEM = Standard Error of the Mean

No statistical analysis was performed on the Biosample cohort females on Postnatal Day 91-98 because there was only one cage in the control group (N=1).

One Gestation Day 15-18 F0 Female feed weight from the 1000 ppm group was excluded as an outlier, along with the relative feed weight for this interval.

Three F0 females from each dose group were removed on GD 18 for biological sampling.

Three F0 females from the 30 and 100 ppm dose groups were removed on LD 4 for biological sampling.

Feed consumption is reported through Postnatal Day 133 for the Neurobehavioral Cohort, and Postnatal Day 119 for the Subchronic Cohort, after which euthanasia of animals across staggered terminal dates occurred.

**** END OF REPORT ****