**Experiment Number:** K10265

Route: Gavage, IV

Species/Strain: Rat/F344/N

# Toxicokinetics Data Summary Test Compound: Methyleugenol

**CAS Number:** 93-15-2

Date Report Requested: 11/09/2016 Time Report Requested: 14:03:57

Lab: Battelle Northwest Laboratory

Male						
	Treatment Groups (mg/kg)					
	37 a	75 a	150 a	37 IV <sup>b</sup>		
	Plasma					
Comin(pred) (ug/mL)				20.0 ± 2.0		
C <sub>max</sub> (ug/mL) *	$0.314 \pm 0.180$	2.22 ± 1.27	1.93 ± 1.12			
T <sub>max</sub> (minute)	5	5	15			
Alpha (min^-1)				0.0405 ± 0.0029		
t <sub>1/2(Alpha)</sub> (minute)				17.1 ± 1.2		
Beta (min^-1)				$0.00530 \pm 0.0007$		
t <sub>1/2(Beta)</sub> (minute)				131 ± 19		
Cl (mL/min/kg)				62.3 ± 1.6		
Cl <sub>1(F)</sub> (mL/min/kg)	1650 ± 160	603 ± 59	321 ± 21			
V1 (L/kg)				11.8 ± 1.7		
AUC <sub>0-t</sub> (ug/mL*min)	21.0 ± 2.2	122 ± 12.0	465 ± 31.0	542.0 ± 15.0		
AUC <sub>inf</sub> (ug/mL*min)				560 ± 35		
F (percent)	3.8					

**Experiment Number:** K10265

Route: Gavage, IV

Species/Strain: Rat/F344/N

# Toxicokinetics Data Summary Test Compound: Methyleugenol CAS Number: 93-15-2

Date Report Requested: 11/09/2016 Time Report Requested: 14:03:57

ber: 93-15-2 Lab: Battelle Northwest Laboratory

Female Female						
	Treatment Groups (mg/kg)					
	37 a	75 ª	150°	37 IV <sup>b</sup>		
	Plasma					
Comin(pred) (ug/mL)				28.9 ± 2.3		
C <sub>max</sub> (ug/mL) *	0.341 ± 0.049	$0.895 \pm 0.067$	4.77 ± 0.81			
Г <sub>тах</sub> (minute)	10	10	5			
Alpha (min^-1)				0.0636 ± 0.0040		
1/2(Alpha) (minute)				10.9 ± 0.7		
Beta (min^-1)				$0.00583 \pm 0.00042$		
1/2(Beta) (minute)				119 ± 9		
CI (mL/min/kg)				59.3 ± 1.0		
Cl <sub>1(F)</sub> (mL/min/kg)	1460 ± 210	885 ± 64	293 ± 19			
√₁ (L/kg)				10.2 ± 0.8		
AUC₀-t (ug/mL*min)	23.0 ± 3.5	82.8 ± 6.1	498 ± 32.0	557.0 ± 10.0		
AUC <sub>inf</sub> (ug/mL*min)				590 ± 26		
F (percent)	4.1					

**Experiment Number:** K10265

Route: Gavage, IV

Species/Strain: Rat/F344/N

Toxicokinetics Data Summary
Test Compound: Methyleugenol
CAS Number: 93-15-2

Date Report Requested: 11/09/2016 Time Report Requested: 14:03:57 Lab: Battelle Northwest Laboratory

## **LEGEND**

Data are displayed as mean ± SEM

\* Data are displayed as mean ± SD

### MODELING METHOD & BEST FIT MODEL

<sup>a</sup> Manual; Plasma MEG concentration-versus-time profiles for both species were characterized by an early absorption phase followed by at least one secondary peak which prevented estimation of elimination rates. No modeling was done on the oral gavage data. Model-independent toxicokinetic parameters were calculated from the gavage data using observed values.

#### **ANALYTE**

Methyleugenol

#### TK PARAMETERS

 $C_{0min(pred)}$  = Fitted plasma concentration at time zero (IV only)

C<sub>max</sub> = Observed or Predicted Maximum plasma (or tissue) concentration

 $T_{max}$  = Time at which  $C_{max}$  predicted or observed occurs

Alpha = Hybrid rate constant of the alpha phase

 $t_{\%(alpha)}$  = Half-life for the alpha phase

Beta = Hybrid rate constant of the beta phase

 $t_{\frac{1}{2}(beta)}$  = Half-life for the beta phase

CI = Clearance, includes total clearance

 $Cl_{1(F)}$  = Apparent clearance of the central compartment, also  $Cl_{(F)}$  for gavage groups in non-compartmental model

 $V_1$  = Volume of distribution of the central compartment, includes  $V_d$  and  $V_{volume}$  of distribution,  $V_z$  apparent volume of distribution NCA,  $V_{app}$  apparent volume of distribution for intravenous studies

AUC<sub>0-t</sub> = Area under the plasma concentration versus time curve, AUC, from time t<sub>i</sub> (initial) to t<sub>f</sub> (final), AUC<sub>last</sub>

AUC<sub>inf</sub> = Area under the plasma concentration versus time curve, AUC, extrapolated to time equals infinity

F = Bioavailability, absolute bioavailability

\*\* END OF REPORT \*\*

<sup>&</sup>lt;sup>b</sup> Nonlinear least-squares fitting program (SAS PROC NLIN; SAS Institute, Inc., Cary, NC); TK parameters of intravenously administered MEG were determined for both species by fitting data to the equation C(t) = Aoe^-alpha\*t + Boe^-beta\*t where C(t) is the plasma MEG concentration at any post-administration time (t), alpha and beta are the hybrid elimination rate constants (min^-1) obtained from the fit, Ao and Bo are the intercepts on the ordinate (concentration) axis of the extrapolated initial and terminal phases, respectively, using a weighting factor of [mean plasma MEG concentration]-2 for rats.